



Extremal conformal structures on projective surfaces

THOMAS METTLER

ABSTRACT. We introduce a new functional \mathcal{E}_p on the space of conformal structures on an oriented projective manifold (M, p) . The non-negative quantity $\mathcal{E}_p([g])$ measures how much p deviates from being defined by a $[g]$ -conformal connection. In the case of a projective surface (Σ, p) , we canonically construct an indefinite Kähler–Einstein structure (h_p, Ω_p) on the total space Y of a fibre bundle over Σ and show that a conformal structure $[g]$ is a critical point for \mathcal{E}_p if and only if a certain lift $\tilde{[g]} : (\Sigma, [g]) \rightarrow (Y, h_p)$ is weakly conformal. In fact, in the compact case $\mathcal{E}_p([g])$ is – up to a topological constant – just the Dirichlet energy of $\tilde{[g]}$. As an application, we prove a novel characterisation of properly convex projective structures among all flat projective structures. As a by-product, we obtain a Gauss–Bonnet type identity for oriented projective surfaces.

CONTENTS

1. Introduction	1
2. Projective and conformal structures	4
3. Projective surfaces and associated bundles	11
4. The variational equations	19
5. Existence of critical points	30
Appendix I. A Gauss–Bonnet type identity	35
References	36

1. Introduction

A *projective structure* on an n -manifold M is an equivalence class p of torsion-free connections on the tangent bundle TM , where two connections are called projectively equivalent if they share the same unparametrised geodesics. A manifold M equipped with a projective structure p will be called a *projective manifold*. A *conformal structure* on M is an equivalence class $[g]$ of Riemannian metrics on M , where two metrics are called conformally equivalent if they differ by a scale factor. Naively, one might think of projective and conformal structures as formally similar, since both arise by defining a notion of equivalence on a geometric structure. However, the formal similarity is more substantial. For instance, Kobayashi has shown [24] that both projective – and conformal structures admit a treatment as Cartan geometries with $|1|$ -graded Lie algebras. Here we exploit the fact that

both structures give rise to affine subspaces modelled on $\Omega^1(M)$ of the infinite-dimensional affine space $\mathfrak{A}(M)$ of torsion-free connections on M . Indeed, it is a classical result due to Weyl [44] that two torsion-free connections on TM are projectively equivalent if and only if their difference – thought of as a section of $S^2(T^*M) \otimes TM$ – is pure trace. Consequently, the representative connections of a projective structure \mathfrak{p} on M define an affine subspace $\mathfrak{A}_{\mathfrak{p}}(M)$ which is modelled on $\Omega^1(M)$. Moreover, it follows from Koszul’s identity, that the torsion-free connections preserving a conformal structure $[g]$ on M are of the form

$${}^g\nabla + g \otimes \beta^\# - \beta \otimes \text{Id} - \text{Id} \otimes \beta,$$

with $g \in [g]$, $\beta \in \Omega^1(M)$ and where ${}^g\nabla$ denotes the Levi-Civita connection of g . Hence, the space of torsion-free $[g]$ -conformal connections on TM is an affine subspace $\mathfrak{A}_{[g]}(M)$ modelled on $\Omega^1(M)$ as well. It is an elementary computation to check that if $\mathfrak{A}_{[g]}(M)$ and $\mathfrak{A}_{\mathfrak{p}}(M)$ intersect, then they do so in a unique point. Therefore, we may ask if in general one can distinguish a point in $\mathfrak{A}_{\mathfrak{p}}(M)$ and a point in $\mathfrak{A}_{[g]}(M)$ which are ‘as close as possible’. This is indeed the case. More precisely, we show that the choice of a conformal structure $[g]$ on (M, \mathfrak{p}) determines a 1-form $A_{[g]}$ on M with values in the endomorphisms of TM , as well as a unique $[g]$ -conformal connection ${}^{[g]}\nabla \in \mathfrak{A}_{[g]}(M)$ so that ${}^{[g]}\nabla + A_{[g]} \in \mathfrak{A}_{\mathfrak{p}}(M)$. The 1-form $A_{[g]}$ appeared previously in the work of Matveev & Trautman [35] and may be thought of as the ‘difference’ between \mathfrak{p} and $[g]$. In particular, if M is oriented, we obtain a $\text{Diff}(M)$ -invariant functional

$$\mathcal{F}(\mathfrak{p}, [g]) = \int_M |A_{[g]}|_g^n d\mu_g.$$

Fixing a projective structure \mathfrak{p} on M , we may consider the functional $\mathcal{E}_{\mathfrak{p}} = \mathcal{F}(\mathfrak{p}, \cdot)$, which is a functional on the space $\mathfrak{C}(M)$ of conformal structures on M only. It is natural to study the infimum of $\mathcal{E}_{\mathfrak{p}}$ among all conformal structures on M , and to ask whether there is actually a minimising conformal structure which achieves this infimum. This infimum – which may be considered as a measure of how far \mathfrak{p} deviates from being defined by a conformal connection – is a new global invariant for oriented projective manifolds.

Of particular interest is the case of surfaces where $\mathcal{E}_{\mathfrak{p}}$ is just the square of the L^2 -norm of $A_{[g]}$ taken with respect to $[g]$ and this is the case that we study in detail in this article. It turns out that in the surface case the functional $\mathcal{E}_{\mathfrak{p}}$ also arises from a rather different viewpoint, which simplifies the computation of its variational equations by using the technique of moving frames.

Inspired by the twistorial construction of holomorphic projective structures by Hitchin [19], it was shown in [13], [42] how to construct a ‘twistor space’ for smooth projective structures. The choice of a projective structure \mathfrak{p} on an oriented surface Σ induces a complex structure on the total space of the disk bundle $Z \rightarrow \Sigma$ whose sections are conformal structures on Σ . In this sense, $\mathcal{E}_{\mathfrak{p}}([g])$ can be interpreted as measuring the failure of $[g](\Sigma) \subset Z$ to be a holomorphic curve in Z . We proceed to show that \mathfrak{p} canonically defines an indefinite Kähler-Einstein structure $(h_{\mathfrak{p}}, \Omega_{\mathfrak{p}})$ on a certain submanifold Y of the projectivised holomorphic cotangent bundle $\mathbb{P}(T_{\mathbb{C}}^*Z^{1,0})$ of Z . Moreover, every conformal structure $[g] : \Sigma \rightarrow Z$ admits a lift $\widetilde{[g]} : \Sigma \rightarrow Y$ so that the variational equations can be expressed as follows:

Theorem A. *Let (Σ, \mathfrak{p}) be an oriented projective surface. A conformal structure $[g]$ on Σ is extremal for \mathfrak{p} if and only if $\widetilde{[g]} : (\Sigma, [g]) \rightarrow (Y, h_{\mathfrak{p}})$ is weakly conformal.*

Here we say that $[g]$ is *extremal* for \mathfrak{p} if it is a critical point of $\mathcal{E}_{\mathfrak{p}}$ with respect to compactly supported variations. Moreover, by weakly conformal we mean that there exists a smooth (and possibly vanishing) function f on Σ so that for some – and hence any – representative metric $g \in [g]$, we have $\widetilde{[g]}^* h_{\mathfrak{p}} = fg$. In fact, in the compact case $\mathcal{E}_{\mathfrak{p}}([g])$ is, up to the topological constant $-2\pi\chi(\Sigma)$, just the Dirichlet energy of $\widetilde{[g]}$. As a consequence, we obtain an optimal lower bound:

Theorem B. *Let (Σ, \mathfrak{p}) be a compact oriented projective surface. Then for every conformal structure $[g] : \Sigma \rightarrow Z$ we have*

$$\frac{1}{2} \int_{\Sigma} \mathrm{tr}_g \widetilde{[g]}^* h_{\mathfrak{p}} d\mu_g \geq -2\pi\chi(\Sigma),$$

with equality if and only if \mathfrak{p} is defined by a $[g]$ -conformal connection.

We then turn to the problem of finding non-trivial examples of projective structures for which $\mathcal{E}_{\mathfrak{p}}$ admits extremal conformal structures. The conformal connection $[g]\nabla$ determined by the choice of a conformal structure $[g]$ on (Σ, \mathfrak{p}) may equivalently be thought of as a torsion-free connection φ on the principal $\mathrm{GL}(1, \mathbb{C})$ -bundle of complex linear coframes of $(\Sigma, [g])$. In addition, the 1-form $A_{[g]}$ turns out to be twice the real part of a section α of $K_{\Sigma}^2 \otimes \overline{K}_{\Sigma}^*$, where K_{Σ} denotes the canonical bundle of $(\Sigma, [g])$. We provide another interpretation of the variational equations by proving that $[g]$ is extremal for \mathfrak{p} if and only if the quadratic differential $\nabla_{\varphi}'' \alpha$ vanishes identically. Here ∇_{φ} denotes the connection induced by φ on $K_{\Sigma}^2 \otimes \overline{K}_{\Sigma}^*$ and ∇_{φ}'' its $(0,1)$ -part. Applying the Riemann–Roch theorem, it follows that a projective structure \mathfrak{p} on the 2-sphere S^2 admits an extremal conformal structure if and only if \mathfrak{p} is defined by a conformal connection.

While there are no non-trivial critical points for projective structures on the 2-sphere, the situation is quite different for surfaces with negative Euler characteristic. Indeed, the condition of having a vanishing quadratic differential appeared previously in the projective differential geometry literature. In the celebrated paper “*Lie groups and Teichmüller space*” [21] Hitchin proposed a generalisation of Teichmüller space \mathcal{H}_2 by identifying a connected component \mathcal{H}_n – nowadays called the *Hitchin component* – in the space of conjugacy classes of representations of $\pi_1(\Sigma)$ into $\mathrm{PSL}(n, \mathbb{R})$.¹ Here Σ denotes a compact oriented surface whose genus exceeds one. Using the theory of Higgs bundles [20] and harmonic map techniques, Hitchin showed that the choice of a conformal structure $[g]$ on Σ gives an identification

$$\mathcal{H}_n \simeq \bigoplus_{\ell=2}^n H^0(\Sigma, K_{\Sigma}^{\ell}).$$

Hitchin conjectured that \mathcal{H}_3 is the space of conjugacy classes of monodromy representations of (flat) properly convex projective structures, a fact later confirmed by Choi and Goldman [10] (the geometric interpretation of the Hitchin component for $n > 3$ is a topic of current interest, c.f. [18], [22], [28] for recent results).

¹More generally, representation into a real split simple Lie group.

Teichmüller space being parametrised by holomorphic quadratic differentials, one might ask if there is a unique choice of a conformal structure on Σ , so that \mathcal{H}_3 is parametrised in terms of cubic holomorphic differentials only. This is indeed the case, as was shown independently by Labourie [27] and Loftin [34] (see also [2] and [14] for recent work treating the non-compact case and the case of convex polygons, as well as [30] treating the case of a general real split rank 2 group). Furthermore, the conformal structure $[g]$ making the quadratic differential vanish is the conformal equivalence class of the so-called *Blaschke metric*, which arises by realising the universal cover of a properly convex projective surface as a complete hyperbolic affine 2-sphere, see in particular [34].

Calling a conformal structure $[g]$ on (Σ, \mathfrak{p}) *closed*, if φ induces a flat connection on $\Lambda^2(T^*\Sigma)$, we obtain a novel characterisation of properly convex projective structures among flat projective structures:

Theorem C. *Let (Σ, \mathfrak{p}) be a compact oriented flat projective surface of negative Euler characteristic. Suppose \mathfrak{p} is properly convex, then the conformal equivalence class of the Blaschke metric is closed and extremal for $\mathcal{E}_{\mathfrak{p}}$. Conversely, if $\mathcal{E}_{\mathfrak{p}}$ admits a closed extremal conformal structure $[g]$, then \mathfrak{p} is properly convex and $[g]$ is the conformal equivalence class of the Blaschke metric of \mathfrak{p} .*

We conclude with some remarks about the possible relation between our functional and the energy functional on Teichmüller space [12], [29] which one can associate to a representation in the Hitchin component. Finally, as a by-product of our ideas, we obtain a Gauss–Bonnet type identity for oriented projective surfaces, which we briefly discuss in Section I.

Acknowledgements

I would like to thank Nigel Hitchin, Gabriel Paternain, Maciej Dunajski, Charles Frances, Karin Melnick and Stefan Rosemann for helpful conversations or correspondence regarding the topic of this article. The author is also grateful to the anonymous referee for helpful remarks. A part of the research for this article was carried out while the author was visiting the Mathematical Institute at the University of Oxford on a postdoctoral fellowship of the Swiss NSF PA00P2_142053. The author would like to thank the Mathematical Institute for its hospitality.

2. Projective and conformal structures

2.1. Preliminaries

Throughout the article, all manifolds are assumed to be connected, have empty boundary and unless stated otherwise, all manifolds and maps are assumed to be smooth, i.e., C^∞ . Also, we adhere to the convention of summing over repeated indices.

2.1.1. Notation

For $\mathbb{F} = \mathbb{R}, \mathbb{C}$ the field of real or complex numbers, we denote by \mathbb{F}^n the space of column vectors of height n and by \mathbb{F}_n the space of row vectors of length n

whose entries are elements of \mathbb{F} . Also, we denote by $\mathbb{FP}^2 = (\mathbb{F}^3 \setminus \{0\})/\mathbb{F}^*$ the space of one-dimensional linear subspaces in \mathbb{F}^3 , that is, the real or complex projective plane. We denote by $\mathbb{S}^2 = (\mathbb{R}^3 \setminus \{0\})/\mathbb{R}^+$ the space of oriented one-dimensional linear subspaces in \mathbb{R}^3 , that is, the projective 2-sphere. Likewise, we write $\mathbb{FP}_2 = (\mathbb{F}_3 \setminus \{0\})/\mathbb{F}^*$ for the dual (real or complex) projective plane and $\mathbb{S}_2 = (\mathbb{R}_3 \setminus \{0\})/\mathbb{R}^+$ for the dual projective 2-sphere. For a non-zero vector $x \in \mathbb{F}^3$ we write $[x]$ for its corresponding point in \mathbb{FP}^2 and for a non-zero vector $\xi \in \mathbb{F}_3$ we write $[\xi]$ for its corresponding point in \mathbb{FP}_2 . For non-zero vectors $x \in \mathbb{R}^3$ and $\xi \in \mathbb{R}_3$ we also use the notation $[x]_+$ and $[\xi]_+$ to denote the corresponding points in \mathbb{S}^2 and \mathbb{S}_2 . Finally, we use the notation $F(\mathbb{F}_3)$ to denote the space of complete flags in \mathbb{F}_3 whose points are pairs (ℓ, Π) with Π being an \mathbb{F} two-dimensional linear subspace of \mathbb{F}_3 containing the line ℓ .

2.1.2. The coframe bundle

Recall that the *coframe bundle* of an n -manifold M is the bundle $\nu : F(T^*M) \rightarrow M$ whose fibre at a point $p \in M$ consists of the linear isomorphisms $u : T_p M \rightarrow \mathbb{R}^n$. The group $\text{GL}(n, \mathbb{R})$ acts transitively from the right on each ν -fibre by the rule $R_a(u) = u \cdot a = a^{-1} \circ u$ for all $a \in \text{GL}(n, \mathbb{R})$. This action turns $\nu : F(T^*M) \rightarrow M$ into a principal right $\text{GL}(n, \mathbb{R})$ -bundle. The coframe bundle is equipped with a tautological \mathbb{R}^n -valued 1-form $\omega = (\omega^i)$ defined by $\omega_u = u \circ \nu'_u$. Note that ω satisfies the equivariance property $R_a^* \omega = a^{-1} \omega$ for all $a \in \text{GL}(n, \mathbb{R})$. The exterior derivative of local coordinates $x : U \rightarrow \mathbb{R}^n$ on M defines a natural section $\tilde{x} : U \rightarrow F(T^*M)$ having the reproducing property $\tilde{x}^* \omega = dx$. We will henceforth write F instead of $F(T^*M)$ whenever M is clear from the context.

2.1.3. Associated bundles

Throughout the article we will frequently make use of the notion of an associated bundle of a principal bundle. The reader will recall that if $\pi : P \rightarrow M$ is a principal right G -bundle and (ρ, N) a pair consisting of a manifold N and a homomorphism $\rho : G \rightarrow \text{Diff}(N)$ into the diffeomorphism group of N , then we obtain an associated fibre bundle with typical fibre N and structure group G whose total space is $P \times_\rho N$, that is, the elements of $P \times_\rho N$ are pairs (u, p) subject to the equivalence relation

$$(u_1, p_1) \sim (u_2, p_2) \iff u_2 = u_1 \cdot g, \quad p_2 = \rho(g^{-1})(p_1), \quad g \in G.$$

A section s of $P \times_\rho N$ is then given by a map $\sigma_s : P \rightarrow N$ which is equivariant with respect to the G -right action on P and the right action of G on N induced by ρ . We say that s is *represented by* σ_s . If N is an affine/linear space and the G -action induced by ρ is affine/linear, then the associated bundle is an affine/vector bundle.

2.2. Projective structures

Recall that the set $\mathfrak{A}(M)$ of torsion-free connections on the tangent bundle of an n -manifold M is the space of sections of an affine bundle $A(M) \rightarrow M$ of rank $\frac{1}{2}n^2(n+1)$ which is modelled on the vector bundle $V = S^2(T^*M) \otimes TM$. We have a canonical trace mapping $\text{tr} : V \rightarrow T^*M$ as well as an inclusion

$$\iota : T^*M \rightarrow V, \quad \nu \mapsto \nu \otimes \text{Id} + \text{Id} \otimes \nu.$$

For every $v \in V$ we let v_0 denote its trace-free part, so that

$$v_0 = v - \frac{1}{(n+1)} \iota(\text{tr } v).$$

A projective structure \mathfrak{p} on a manifold M of dimension $n > 1$ is an equivalence class of torsion-free connections on TM , where two connections are declared to be equivalent if they share the same unparametrised geodesics. Weyl [44] observed the following:

Lemma 2.1. *Two torsion-free connections ∇ and ∇' on TM are projectively equivalent if and only if $(\nabla - \nabla')_0 = 0$.*

Consequently, the set $\mathfrak{P}(M)$ of projective structures on M is the space of sections of an affine bundle $\mathbf{P}(M) \rightarrow M$ of rank $\frac{1}{2}(n+2)n(n-1)$ which is modelled on the traceless part V_0 of the vector bundle V . We will use the notation $\mathfrak{p}(\nabla)$ for the projective structure \mathfrak{p} that is defined by a connection ∇ . A consequence of Weyl's result is that the set of representative connections of a projective structure \mathfrak{p} is an affine subspace $\mathfrak{A}_{\mathfrak{p}}(M) \subset \mathfrak{A}(M)$ of the space of torsion-free connections which is modelled on the space of 1-forms on M .

2.3. Conformal structures

A conformal structure on a manifold M of dimension $n > 1$ is an equivalence class $[g]$ of Riemannian metrics on M , where two metrics g and \hat{g} are declared to be equivalent if there exists a smooth function f on M so that $\hat{g} = e^{2f}g$. Equivalently, a conformal structure $[g]$ on M is a (smooth) choice of a coframe for every point p in M , well defined up to orthogonal transformation and scaling. Consequently, the set $\mathfrak{C}(M)$ of conformal structures on M is the space of sections of $\mathbf{C}(M) = F/(\mathbb{R}^+ \times \text{O}(n)) \rightarrow M$, where $\mathbb{R}^+ \times \text{O}(n)$ is the subgroup of $\text{GL}(n, \mathbb{R})$ consisting of matrices a having the property that aa^t is a non-zero multiple of the identity matrix.

A torsion-free connection ∇ on TM is called a *Weyl connection* or *conformal connection* for the conformal structure $[g]$ on M if the parallel transport maps of ∇ are angle-preserving with respect to $[g]$. A torsion-free connection ∇ is $[g]$ -conformal if for some (and hence any) representative metric $g \in [g]$ there exists a 1-form β on M such that

$$\nabla g = 2\beta \otimes g.$$

It is a simple consequence of Koszul's identity that the $[g]$ -conformal connections are of the form

$$(2.1) \quad (g, \beta)\nabla = {}^g\nabla + g \otimes \beta^\sharp - \beta \otimes \text{Id} - \text{Id} \otimes \beta,$$

where $g \in [g]$, β is a 1-form on M with g -dual vector field β^\sharp and ${}^g\nabla$ denotes the Levi-Civita connection of g . Consequently, the set of $[g]$ -conformal connections defines an affine subspace $\mathfrak{A}_{[g]}(M) \subset \mathfrak{A}(M)$ which is modelled on the space of 1-forms on M as well. For later usage we also record that for every smooth function f on M we have

$$(\exp(2f)g, \beta + df)\nabla = (g, \beta)\nabla,$$

as the reader may easily verify using the identity [3, Theorem 1.159]

$$(2.2) \quad \exp(2f)g \nabla = g \nabla - g \otimes g \nabla f + \iota(df).$$

In particular, if β is exact, so that $\beta = df$ for some smooth function f on M , then $(g, \beta) \nabla = \exp(-2f)g \nabla$ and hence the conformal connection determined by (g, β) is the Levi-Civita connection of the metric $e^{-2f}g$.

We also use the notation $[g] \nabla$ for a connection preserving the conformal structure $[g]$.

2.4. Compatibility of projective and conformal structures

Since both projective – and conformal structures give rise to affine subspaces of $\mathfrak{A}(M)$ of the same type, we may ask how two such spaces intersect.

Lemma 2.2. *Let $[g]$ be a conformal – and \mathfrak{p} a projective structure on M . Then $\mathfrak{A}_{[g]}(M)$ and $\mathfrak{A}_{\mathfrak{p}}(M)$ intersect in at most one point.*

Proof. Suppose the $[g]$ -conformal connections $[g] \nabla$ and $[g] \hat{\nabla}$ are both elements in $\mathfrak{A}_{\mathfrak{p}}(M)$. Then, by Theorem 2.1, there exists a 1-form Υ on M so that

$$[g] \nabla = [g] \hat{\nabla} + \iota(\Upsilon).$$

Fixing a Riemannian metric g defining $[g]$, we also have 1-forms $\beta, \hat{\beta}$ on M so that

$$[g] \nabla = g \nabla + g \otimes \beta^{\#} - \iota(\beta) \quad \text{and} \quad [g] \hat{\nabla} = g \nabla + g \otimes \hat{\beta}^{\#} - \iota(\hat{\beta}).$$

Applying these formulae we obtain

$$\iota(\Upsilon + \beta - \hat{\beta}) = g \otimes (\beta^{\#} - \hat{\beta}^{\#}).$$

Taking the trace gives

$$(n+1)(\Upsilon + \beta - \hat{\beta}) = \beta - \hat{\beta},$$

so that $\Upsilon = -\frac{n}{(n+1)}(\beta - \hat{\beta})$. Therefore we must have

$$\iota(\beta - \hat{\beta}) = (n+1)g \otimes (\beta^{\#} - \hat{\beta}^{\#}).$$

Contracting this last equation with the dual metric $g^{\#}$ implies

$$0 = (n+2)(n-1)(\beta^{\#} - \hat{\beta}^{\#}),$$

so that $\beta = \hat{\beta}$ provided $n > 1$. It follows that Υ vanishes too, therefore $[g] \nabla = [g] \hat{\nabla}$, as claimed. \square

Remark 2.3. Theorem 2.2 raises the question whether or not one can still determine a unique point $[g] \nabla \in \mathfrak{A}_{[g]}(M)$ and a unique point $\nabla \in \mathfrak{A}_{\mathfrak{p}}(M)$ in the general case, where $\mathfrak{A}_{[g]}(M)$ and $\mathfrak{A}_{\mathfrak{p}}(M)$ might not intersect. Formally speaking, we are interested in maps

$$\psi = (\psi^1, \psi^2) : \mathfrak{P}(M) \times \mathfrak{C}(M) \rightarrow \mathfrak{A}(M) \times \mathfrak{A}(M)$$

satisfying the following properties:

- (i) $\psi^1(\mathfrak{p}, [g]) \in \mathfrak{A}_{\mathfrak{p}}(M)$ and $\psi^2(\mathfrak{p}, [g]) \in \mathfrak{A}_{[g]}(M)$;
- (ii) If $\mathfrak{A}_{\mathfrak{p}}(M) \cap \mathfrak{A}_{[g]}(M)$ is non-empty, then $\psi^2(\mathfrak{p}, [g]) - \psi^1(\mathfrak{p}, [g]) = 0$;

- (iii) ψ is equivariant with respect to the natural right action of the diffeomorphism group $\text{Diff}(M)$ on $\mathfrak{P}(M) \times \mathfrak{C}(M)$ and $\mathfrak{A}(M) \times \mathfrak{A}(M)$.

We will next discuss a geometrically natural and explicit map ψ having these properties.

To this end let g be a Riemannian metric on M and ∇ a torsion-free connection on TM . Consider the first-order differential operator for g mapping into the space of 1-forms on M with values in $\text{End}(TM)$

$$(2.3) \quad g \mapsto A_{[g]} = (\nabla - {}^g\nabla - g \otimes X_g)_0,$$

where $X_g \in \Gamma(TM)$ is

$$(2.4) \quad X_g = \frac{(n+1)}{(n+2)(n-1)} \text{tr} \left(g^\# \otimes (\nabla - {}^g\nabla)_0 \right).$$

The following result is essentially contained in [35] – except for (vi). For the convenience of the reader we include a proof.

Theorem 2.4 (Matveev & Trautman, [35]). *The 1-form $A_{[g]}$ has the following properties:*

- (i) *the endomorphism $A_{[g]}(X)$ is trace-free for all $X \in \Gamma(TM)$;*
- (ii) *for all $X, Y \in \Gamma(TM)$ we have $A_{[g]}(X)Y = A_{[g]}(Y)X$;*
- (iii) *$A_{[g]}$ only depends on the projective equivalence class of ∇ ;*
- (iv) *$A_{[g]}$ only depends on the conformal equivalence class of g ;*
- (v) *$A_{[g]} \equiv 0$ if and only if there exists a $[g]$ -conformal connection which is projectively equivalent to ∇ ;*
- (vi) *for $n = 2$ the endomorphism $A_{[g]}(X)$ is symmetric with respect to $[g]$ for all $X \in \Gamma(TM)$;*

Proof. The properties (i) and (ii) are obvious from the definition.

(iii) Recall that two affine torsion-free connections ∇ and $\hat{\nabla}$ are projectively equivalent if and only if $(\nabla - \hat{\nabla})_0 = 0$. The claim follows from the linearity of the “taking the trace-free part” operation.

(iv) Let $\hat{g} = e^{2f}g$ for some smooth real-valued function f on M . Then we have

$$\hat{g}\nabla = {}^g\nabla - g \otimes {}^g\nabla f + \iota(df)$$

and hence

$$\begin{aligned} (\nabla - \hat{g}\nabla)_0 &= (\nabla - {}^g\nabla)_0 + (g \otimes {}^g\nabla f - \iota(df))_0 \\ &= (\nabla - {}^g\nabla)_0 + (g \otimes {}^g\nabla f)_0 \\ &= (\nabla - {}^g\nabla)_0 + g \otimes {}^g\nabla f - \frac{1}{(n+1)}\iota(df). \end{aligned}$$

We obtain

$$\begin{aligned} X_{\hat{g}} &= \frac{(n+1)}{(n+2)(n-1)} \text{tr} \left[\hat{g}^\# \otimes \left((\nabla - {}^g\nabla)_0 + g \otimes {}^g\nabla f - \frac{1}{(n+1)}\iota(df) \right) \right] \\ &= e^{-2f} \left(X_g + \frac{n(n+1)}{(n+2)(n-1)} {}^g\nabla f - \frac{2}{(n+2)(n-1)} {}^g\nabla f \right) \\ &= e^{-2f} (X_g + {}^g\nabla f). \end{aligned}$$

This gives

$$\begin{aligned}\hat{g}\nabla + \hat{g} \otimes X_{\hat{g}} &= {}^g\nabla - g \otimes {}^g\nabla f + \iota(df) + e^{2f} g \otimes e^{-2f} (X_g + {}^g\nabla f) \\ &= {}^g\nabla + g \otimes X_g + \iota(df),\end{aligned}$$

so that

$$\left(\hat{g}\nabla + \hat{g} \otimes X_{\hat{g}}\right)_0 = \left({}^g\nabla + g \otimes X_g\right)_0,$$

which shows that $A_{[g]}$ does indeed only depend on the conformal class of g .

(v) Recall that the $[g]$ -conformal connections are of the form

$$[g]\nabla = {}^g\nabla + g \otimes \beta^\# - \iota(\beta),$$

where g is any metric in the conformal class $[g]$ and β is some 1-form on M . Therefore we have

$$\left([g]\nabla - {}^g\nabla\right)_0 = \left(g \otimes \beta^\#\right)_0 = g \otimes \beta^\# - \frac{1}{(n+1)}\iota(\beta)$$

and thus as before we compute that $X_g = \beta^\#$. We obtain

$$\begin{aligned}A_{[g]} &= \left[[g]\nabla - \left({}^g\nabla + g \otimes X_g\right)\right]_0 \\ &= \left[{}^g\nabla + g \otimes \beta^\# - \iota(\beta) - {}^g\nabla - g \otimes \beta^\#\right]_0 = [-\iota(\beta)]_0 = 0.\end{aligned}$$

Conversely, suppose \mathfrak{p} is a projective structure for which there exists a conformal structure $[g]$ with $A_{[g]} \equiv 0$. Fixing a Riemannian metric $g \in [g]$ and a \mathfrak{p} -representative connection ∇ , we must have

$$\nabla - ({}^g\nabla + g \otimes X_g) = \iota(\beta),$$

for some 1-form β on M . Adding $\iota((X_g)^\flat)$ gives

$$\nabla - \left({}^g\nabla + g \otimes X_g - \iota((X_g)^\flat)\right) = \iota(\beta + (X_g)^\flat),$$

so that Theorem 2.1 implies that ∇ and the $[g]$ -conformal connection

$${}^g\nabla + g \otimes X_g - \iota((X_g)^\flat)$$

are projectively equivalent.

(vi) Let now $n = 2$. We need to show that for $g \in [g]$ and all vector fields $X, Y, Z \in \Gamma(TM)$, we have

$$g(A_{[g]}(X)Y, Z) = g(Y, A_{[g]}(X)Z).$$

Without losing generality, we can assume that locally $g = (dx^1)^2 + (dx^2)^2$ for coordinates $x = (x^1, x^2) : U \rightarrow \mathbb{R}^2$ on M . Let Γ_{jk}^i denote the Christoffel symbols of ∇ with respect to x . Since the Christoffel symbols of ${}^g\nabla$ vanish identically on U , we obtain with a simple calculation

$$X_g = -\frac{3}{4}(w_1 + w_3)\frac{\partial}{\partial x^1} + \frac{3}{4}(w_0 + w_2)\frac{\partial}{\partial x^2},$$

where

$$w_0 = \Gamma_{11}^2, \quad 3w_1 = -\Gamma_{11}^1 + 2\Gamma_{12}^2, \quad 3w_2 = -2\Gamma_{12}^1 + \Gamma_{22}^2, \quad w_3 = -\Gamma_{22}^1.$$

Likewise, we compute

$$A_{[g]} = \frac{1}{2} (a_1 e^{11}_1 - a_2 e^{11}_2 - a_2 e^{12}_1 - a_1 e^{12}_2 \\ - a_2 e^{21}_1 - a_1 e^{21}_2 - a_1 e^{22}_1 + a_2 e^{22}_2)$$

where we write $e^{ij}_k = dx^i \otimes dx^j \otimes \frac{\partial}{\partial x^k}$ and

$$a_1 = \frac{1}{2}(w_3 - 3w_1), \quad a_2 = \frac{1}{2}(3w_2 - w_0).$$

The claim follows from an elementary calculation. \square

Remark 2.5. By construction, the 1-form $A_{[g]}$ vanishes identically if and only if ∇ is projectively equivalent to a conformal connection. The necessary and sufficient conditions for a torsion-free connection to be projectively equivalent to a Levi-Civita connection were given in [4]. The reader may also consult [7] for the role of Einstein metrics in projective differential geometry.

As a corollary to Theorem 2.4 and Theorem 2.2 we obtain the following result.

Corollary 2.6. *For every conformal structure $[g]$ on the projective manifold (M, \mathfrak{p}) , there exists a unique $[g]$ -conformal connection $^{[g]}\nabla$ so that $^{[g]}\nabla + A_{[g]} \in \mathfrak{p}$.*

Note that Theorem 2.6 provides a unique point $^{[g]}\nabla \in \mathfrak{A}_{[g]}(M)$ and a unique point $^{[g]}\nabla + A_{[g]} \in \mathfrak{A}_{\mathfrak{p}}(M)$. We may define

$$\psi(\mathfrak{p}, [g]) = \left(^{[g]}\nabla + A_{[g]}, ^{[g]}\nabla \right).$$

Since the map which sends a Riemannian metric to its Levi-Civita connection is equivariant with respect to the action of $\text{Diff}(M)$ on the space of Riemannian metrics and on $\mathfrak{A}(M)$, it follows that the map ψ has all the properties listed in Theorem 2.3.

Proof of Theorem 2.6. Let ∇ be a connection defining \mathfrak{p} and g a smooth metric defining $[g]$. Set

$$^{[g]}\nabla = {}^g\nabla + g \otimes X_g - (X_g)^\flat \otimes \text{Id} - \text{Id} \otimes (X_g)^\flat,$$

where X_g is defined as before (see (2.4)). Then, property (i) of $A_{[g]}$ proved in Theorem 2.4 implies that

$$\left(\nabla - \left(^{[g]}\nabla + A_{[g]} \right) \right)_0 = \left(\nabla - ({}^g\nabla + g \otimes X_g) \right)_0 - A_{[g]} = A_{[g]} - A_{[g]} = 0,$$

so that $^{[g]}\nabla + A_{[g]}$ is projectively equivalent to ∇ by Theorem 2.1. If $^{[g]}\nabla'$ is another $[g]$ -conformal connection so that $^{[g]}\nabla' + A_{[g]}$ defines \mathfrak{p} , then

$$\left(^{[g]}\nabla - ^{[g]}\nabla' \right)_0 = 0,$$

hence $^{[g]}\nabla = ^{[g]}\nabla'$ by Theorem 2.2. \square

2.5. A diffeomorphism invariant functional

We will henceforth assume M to be oriented. For a pair $(p, [g])$ consisting of a projective structure and a conformal structure on M , we consider the non-negative n -form $|A_{[g]}|_g^n d\mu_g$, where g is any metric defining $[g]$, the n -form $d\mu_g$ denotes its volume form and where $A_{[g]}$ is computed with respect to p . For $f \in C^\infty(M)$ we have

$$|A_{[g]}|_{e^{2f}g} = e^{-f} |A_{[g]}|_g \quad \text{and} \quad d\mu_{e^{2f}g} = e^{nf} d\mu_g,$$

it follows that $|A_{[g]}|_g^n d\mu_g$ depends only on the conformal structure $[g]$. Consequently, we obtain a non-negative functional

$$\mathcal{F} : \mathfrak{P}(M) \times \mathfrak{C}(M) \rightarrow \mathbb{R}_0^+ \cup \{\infty\}, \quad (p, [g]) \mapsto \int_M |A_{[g]}|_g^n d\mu_g.$$

By construction, \mathcal{F} is invariant under simultaneous action of $\text{Diff}(M)$ on $\mathfrak{P}(M)$ and $\mathfrak{C}(M)$.

We may also fix a projective structure p on M and define $\mathcal{E}_p = \mathcal{F}[(p, \cdot)]$ which is a functional on $\mathfrak{C}(M)$ only. We may study the infimum of \mathcal{E}_p among all conformal structures on M , and ask whether there is actually a minimising conformal structure which achieves this infimum. The infimum

$$\Gamma\delta(M, p) := \inf_{[g] \in \mathfrak{C}(M)} \mathcal{E}_p([g]),$$

which may be considered as a measure of how far p deviates from being defined by a conformal connection, is a new global invariant for oriented projective manifolds. Note that reversing the role of p and $[g]$ does not give us a global invariant for conformal manifolds. Clearly, fixing a conformal structure and considering the infimum over $\mathfrak{P}(M)$ yields zero for every choice of conformal structure $[g]$.

3. Projective surfaces and associated bundles

A natural case to consider is $n = 2$, where \mathcal{F} is just the square of the L^2 -norm of $A_{[g]}$ taken with respect to $[g]$. We will henceforth consider the surface case only.

There are several natural geometric spaces fibering over an oriented projective surface which we will discuss next. Before doing so, we recall a result of Cartan [9], which canonically associates a principal bundle together with a “connection” to every projective manifold. The reader interested in a description of Cartan’s construction using modern language may also consult [26]. For additional background on Cartan geometries the reader may also consult [8].

3.1. Cartan’s normal projective connection

Let Σ be an oriented surface and let $G \simeq \mathbb{R}_2 \rtimes \text{GL}^+(2, \mathbb{R})$ denote the two-dimensional orientation preserving affine group which we think of as the subgroup of $\text{SL}(3, \mathbb{R})$ consisting of matrices of the form

$$b \rtimes a = \begin{pmatrix} \det a^{-1} & b \\ 0 & a \end{pmatrix},$$

for $b \in \mathbb{R}_2$ and $a \in \text{GL}^+(2, \mathbb{R})$. We denote by $\nu : F^+ \rightarrow \Sigma$ the principal right $\text{GL}^+(2, \mathbb{R})$ -bundle of coframes that are orientation preserving with respect to the

chosen orientation on Σ and the standard orientation on \mathbb{R}^2 . We define a right G -action on $F^+ \times \mathbb{R}_2$ by the rule

$$(3.1) \quad (u, \xi) \cdot (b \rtimes a) = (\det a^{-1} a^{-1} \circ u, \xi a \det a + b \det a),$$

for all $b \rtimes a \in G$. Here $\xi : F^+ \times \mathbb{R}_2 \rightarrow \mathbb{R}_2$ denotes the projection onto the latter factor. This action turns $\pi : F^+ \times \mathbb{R}_2 \rightarrow \Sigma$ into a principal right G -bundle over Σ , where $\pi : F^+ \times \mathbb{R}_2 \rightarrow \Sigma$ denotes the natural basepoint projection. Suppose ∇ is a torsion-free connection on $T\Sigma$ with connection 1-form $\eta = (\eta_j^i)$ on F^+ so that we have the structure equations²

$$\begin{aligned} d\omega^i &= -\eta_j^i \wedge \omega^j, \\ d\eta_j^i &= -\eta_k^i \wedge \eta_j^k + (\delta_{[k}^i S_{l]j} - S_{[kl]j} \delta_j^i) \omega^k \wedge \omega^l, \end{aligned}$$

where $S = (S_{ij})$ represents the *projective Schouten tensor* $\text{Schout}(\nabla)$ of ∇ and ω^i the components of the tautological \mathbb{R}^n -valued 1-form ω on F . Recall that the Schouten tensor is defined as

$$(3.2) \quad \text{Schout}(\nabla) = \text{Ric}^+(\nabla) - \frac{1}{3} \text{Ric}^-(\nabla),$$

where $\text{Ric}^\pm(\nabla)$ denote the symmetric and anti-symmetric part of the Ricci curvature of ∇ . On $P = F^+ \times \mathbb{R}_2$ we define the $\mathfrak{sl}(3, \mathbb{R})$ -valued 1-form

$$(3.3) \quad \theta = \begin{pmatrix} -\frac{1}{3} \text{tr } \eta - \xi \omega & d\xi - \xi \eta - (S\omega)^t - \xi \omega \xi \\ \omega & \eta - \frac{1}{3} \text{I tr } \eta + \omega \xi \end{pmatrix}.$$

The reader may check that the pair $(\pi : P \rightarrow \Sigma, \theta)$ defines a *Cartan geometry* of type $(\text{SL}(3, \mathbb{R}), G)$, that is, $\pi : P \rightarrow \Sigma$ is a principal right G -bundle and θ is an $\mathfrak{sl}(3, \mathbb{R})$ -valued 1-form on P satisfying the following properties:

- (i) $\theta_u : T_u P \rightarrow \mathfrak{sl}(3, \mathbb{R})$ is an isomorphism for every $u \in P$;
- (ii) $(R_g)^* \theta = g^{-1} \theta g$ for every $g \in G$;
- (iii) $\theta(X_v) = v$ for every fundamental vector field X_v generated by an element v in the Lie algebra of G .

Moreover, writing $\theta = (\theta_j^i)_{i,j=0,1,2}$, the Cartan geometry $(\pi : P \rightarrow \Sigma, \theta)$ also satisfies:

- (iv) for every non-zero $x \in \mathbb{R}^2$, the integral curves of the vector field X_x defined by the equations

$$\theta(X_x) = \begin{pmatrix} 0 & 0 \\ x & 0 \end{pmatrix}$$

project to Σ to become geodesics of \mathfrak{p} and conversely all geodesics of \mathfrak{p} arise in this way;

- (v) the π -pullback of an orientation compatible volume form on Σ is a positive multiple of $\theta_0^1 \wedge \theta_0^2$;

²Indices in round brackets are symmetrised over and indices in square brackets are anti-symmetrised over, for instance, we write $S_{(ij)} = \frac{1}{2} (S_{ij} + S_{ji})$ and $S_{[ij]} = \frac{1}{2} (S_{ij} - S_{ji})$ so that $S_{ij} = S_{(ij)} + S_{[ij]}$.

(vi) the curvature 2-form $\Theta = d\theta + \theta \wedge \theta$ is

$$(3.4) \quad \Theta = d\theta + \theta \wedge \theta = \begin{pmatrix} 0 & L_1\theta_0^1 \wedge \theta_0^2 & L_2\theta_0^1 \wedge \theta_0^2 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

for unique curvature functions $L_i : P \rightarrow \mathbb{R}$.

Remark 3.1. Cartan's bundle is unique in the following sense: If $(\hat{\pi} : \hat{P} \rightarrow \Sigma, \hat{\theta})$ is another Cartan geometry of type $(\mathrm{SL}(3, \mathbb{R}), G)$ so that the properties (iv), (v) and (vi) hold, then there exists a G -bundle isomorphism $\psi : P \rightarrow \hat{P}$ satisfying $\psi^*\hat{\theta} = \theta$.

A projective structure \mathfrak{p} on Σ is called *flat* if every point $p \in \Sigma$ has a neighbourhood U_p which is diffeomorphic to a subset of \mathbb{RP}^2 in such a way that the geodesics of \mathfrak{p} contained in U_p are mapped onto (segments) of projective lines $\mathbb{RP}^1 \subset \mathbb{RP}^2$. Furthermore, a torsion-free connection ∇ on $T\Sigma$ is called *projectively flat* if $\mathfrak{p}(\nabla)$ is flat. Using Cartan's connection, one can show that a projective structure \mathfrak{p} is flat if and only if the functions L_1 and L_2 vanish identically. Another consequence of Cartan's result is that there exists a unique 1-form $\lambda \in \Omega^1(\Sigma, \Lambda^2(T^*\Sigma))$ so that

$$\pi^*\lambda = (L_1\theta_0^1 + L_2\theta_0^2) \otimes \theta_0^1 \wedge \theta_0^2.$$

The 1-form λ was first discovered by R. Liouville [32], hence we call λ the *Liouville curvature* of \mathfrak{p} . In particular, the Liouville curvature is the complete obstruction to flatness of a two-dimensional projective structure.

Example 3.2. Note that the left action of $\mathrm{SL}(3, \mathbb{R})$ on \mathbb{R}^3 by matrix multiplication descends to define a transitive left action on the projective 2-sphere \mathbb{S}^2 . The stabiliser subgroup of the element $[(1 \ 0 \ 0)^t]$ is the group $G \subset \mathrm{SL}(3, \mathbb{R})$ so that $\mathbb{S}^2 \simeq \mathrm{SL}(3, \mathbb{R})/G$. Taking θ to be the Maurer-Cartan form of $\mathrm{SL}(3, \mathbb{R})$, the pair $(\pi : \mathrm{SL}(3, \mathbb{R}) \rightarrow \mathbb{S}^2, \theta)$ is a Cartan geometry of type $(\mathrm{SL}(3, \mathbb{R}), G)$ defining an orientation and projective structure $\mathfrak{p}_{\mathrm{can}}$ on \mathbb{S}^2 whose geodesics are the “great circles”. Since $d\theta + \theta \wedge \theta = 0$, this projective structure is flat. We call $\mathfrak{p}_{\mathrm{can}}$ the *canonical flat projective structure* on \mathbb{S}^2 .

3.2. The twistor space

Inspired by Hitchin's twistorial description of holomorphic projective structures on complex surfaces [19], it was shown in [13, 42] how to construct a “twistor space” for smooth projective structures. For what follows it will be convenient to construct the twistor space in the smooth category by using the Cartan geometry of a projective surface.

Let therefore (Σ, \mathfrak{p}) be an oriented projective surface with Cartan geometry $(\pi : P \rightarrow \Sigma, \theta)$. By construction, the quotient of P by the normal subgroup $\mathbb{R}_2 \rtimes \{\mathrm{Id}\} \subset G$ is isomorphic to the bundle $\nu : F^+ \rightarrow \Sigma$ of orientation preserving coframes of Σ . In particular, the choice of a conformal structure $[g]$ on Σ corresponds to a section of the fibre bundle $C(\Sigma) \simeq P/(\mathbb{R}_2 \rtimes \mathrm{CO}(2)) \rightarrow \Sigma$. Here $\mathrm{CO}(2) = \mathbb{R}^+ \times \mathrm{SO}(2)$ is the linear orientation preserving conformal group. By construction, the typical fibre of the bundle $C(\Sigma) \rightarrow \Sigma$ is diffeomorphic to $\mathrm{GL}^+(2, \mathbb{R})/\mathrm{CO}(2) \simeq \mathrm{SL}(2, \mathbb{R})/\mathrm{SO}(2)$, that is, the open unit disk $D^2 \subset \mathbb{C}$.

We write the elements of the group $\mathbb{R}_2 \rtimes \text{CO}(2)$ in the following form

$$z \rtimes r e^{i\phi} = \begin{pmatrix} r^{-2} & \text{Re}(z) & \text{Im}(z) \\ 0 & r \cos \phi & r \sin \phi \\ 0 & -r \sin \phi & r \cos \phi \end{pmatrix}, \quad z \in \mathbb{C}, r e^{i\phi} \in \mathbb{C}^*.$$

Property (iii) of the Cartan geometry $(\pi : P \rightarrow \Sigma, \theta)$ implies that the (real – or complex-valued) 1-forms on P that are semibasic³ for the quotient projection $\mu : P \rightarrow \text{C}(\Sigma)$ are complex linear combinations of the complex-valued 1-forms

$$(3.5) \quad \zeta_1 = \theta_0^1 + i\theta_0^2, \quad \zeta_2 = (\theta_1^1 - \theta_2^2) + i(\theta_1^2 + \theta_2^1)$$

and their complex conjugates. The equivariance property (ii) of the Cartan geometry gives

$$(3.6) \quad (R_{z \rtimes r e^{i\phi}})^* \begin{pmatrix} \zeta_1 \\ \zeta_2 \end{pmatrix} = \begin{pmatrix} \frac{1}{r^3} e^{i\phi} & 0 \\ \frac{z}{r} e^{i\phi} & e^{2i\phi} \end{pmatrix} \begin{pmatrix} \zeta_1 \\ \zeta_2 \end{pmatrix}.$$

It follows that there exists a unique almost complex structure \mathfrak{J} on $\text{C}(\Sigma)$ having the property that a complex-valued 1-form on P is the pullback of a $(1,0)$ -form on $\text{C}(\Sigma)$ if and only if it is a complex linear combination of ζ_1 and ζ_2 . Indeed, locally we may use a section s of the bundle $\mu : P \rightarrow \text{C}(\Sigma)$ to pull down the forms ζ_1, ζ_2 onto the domain of definition $U \subset \text{C}(\Sigma)$ of s . Since ζ_1, ζ_2 are semi-basic for the projection $\mu : P \rightarrow \text{C}(\Sigma)$, it follows that the pulled down forms are linearly independent over \mathbb{C} at each point of U . Hence we obtain a unique almost complex structure \mathfrak{J} on U whose $(1,0)$ -forms are $s^* \zeta_1, s^* \zeta_2$. The equivariance (3.6) implies that \mathfrak{J} is independent of the choice of the section s and extends to all of $\text{C}(\Sigma)$. Using property (vi) of the Cartan geometry the reader may easily verify that

$$d\zeta_1 = d\zeta_2 = 0, \quad \text{mod } \zeta_1, \zeta_2.$$

It follows from the Newlander-Nirenberg theorem that \mathfrak{J} is integrable, thus giving $\text{C}(\Sigma)$ the structure of a complex surface which we will denote by Z and which – abusing language – we call the *twistor space* of the projective surface (Σ, \mathfrak{p}) .

3.3. An indefinite Kähler-Einstein 3-fold

From (3.6) it follows that the holomorphic cotangent bundle $T_{\mathbb{C}}^* Z^{1,0} \rightarrow Z$ is the bundle associated to $\mu : P \rightarrow Z$ via the complex two-dimensional representation $\rho : \mathbb{R}_2 \rtimes \text{CO}(2) \rightarrow \text{GL}(2, \mathbb{C})$ defined by the rule

$$(3.7) \quad \rho(z \rtimes r e^{i\phi})(w_1 \ w_2) = (w_1 \ w_2) \begin{pmatrix} \frac{1}{r^3} e^{i\phi} & 0 \\ \frac{z}{r} e^{i\phi} & e^{2i\phi} \end{pmatrix}$$

for all $(w_1 \ w_2) \in \mathbb{C}_2$. In particular, the form ζ_1 is well defined on Z up to complex-scale and hence may be thought of as a section of the projective holomorphic cotangent bundle $\mathbb{P}(T_{\mathbb{C}}^* Z^{1,0}) \rightarrow Z$. Abusing notation, we write $\zeta_1(Z) \subset \mathbb{P}(T_{\mathbb{C}}^* Z^{1,0})$ to denote the image of Z under this section. We now have:

³Recall that a differential form α is said to be semibasic for the projection $P \rightarrow \text{C}(\Sigma)$ if the interior product $X \lrcorner \alpha$ vanishes for every vector field X tangent to the fibres of $P \rightarrow \text{C}(\Sigma)$.

Lemma 3.3. *There exists a unique integrable almost complex structure on the quotient $P/\mathrm{CO}(2)$ having the property that its $(1,0)$ -forms pull back to P to become linear combinations of the forms*

$$(3.8) \quad \zeta_1 = \theta_0^1 + i\theta_0^2, \quad \zeta_2 = (\theta_1^1 - \theta_2^2) + i(\theta_2^1 + \theta_1^2), \quad \zeta_3 = \theta_1^0 + i\theta_2^0.$$

*Furthermore, with respect to this complex structure $P/\mathrm{CO}(2)$ is biholomorphic to $Y = \mathbb{P}(T_{\mathbb{C}}^*Z^{1,0}) \setminus \zeta_1(Z)$ in such a way that the standard holomorphic contact structure on Y is identified with the subbundle of $T_{\mathbb{C}}(P/\mathrm{CO}(2))^{1,0}$ defined by the equation $\zeta_2 = 0$.*

Proof. Again, it follows from the property (iii) of the Cartan connection θ that the 1-forms that are semibasic for the quotient projection $\tau : P \rightarrow P/\mathrm{CO}(2)$ are linear combinations of the forms $\zeta_1, \zeta_2, \zeta_3$ and their complex conjugates. Here $\mathrm{CO}(2) \subset G$ is the subgroup consisting of elements of the form $0 \rtimes re^{i\phi}$. Writing $re^{i\phi}$ instead of $0 \rtimes re^{i\phi}$ and $\zeta = (\zeta_i)$, we compute from the equivariance property (ii) of θ that we have

$$(3.9) \quad (R_{re^{i\phi}})^* \begin{pmatrix} \zeta_1 \\ \zeta_2 \\ \zeta_3 \end{pmatrix} = \begin{pmatrix} \frac{1}{r^3}e^{i\phi} & 0 & 0 \\ 0 & e^{2i\phi} & 0 \\ 0 & 0 & r^3e^{i\phi} \end{pmatrix} \begin{pmatrix} \zeta_1 \\ \zeta_2 \\ \zeta_3 \end{pmatrix}.$$

It follows as before that there exists a unique almost complex structure \mathfrak{J} on the quotient $P/\mathrm{CO}(2)$ having the property that its $(1,0)$ -forms pull back to P to become linear combinations of the forms $\zeta_1, \zeta_2, \zeta_3$. Suppose there exists a 1-form $\gamma = (\gamma_{ij})$ on P with values in $\mathfrak{gl}(3, \mathbb{C})$, so that $d\zeta = -\gamma \wedge \zeta$, then it follows again from the Newlander–Nirenberg theorem that \mathfrak{J} is integrable. Clearly, if such a γ exists, then it is not unique. Defining $\hat{\gamma} = (\hat{\gamma}_{ij})$, with $\hat{\gamma}_{ij} = \gamma_{ij} + T_{ijk}\zeta_k$ for some complex-valued functions satisfying $T_{ijk} = T_{ikj}$ on P will also work. We can exploit this freedom and make γ take values in the Lie algebra

$$\mathfrak{u}(2, 1) = \left\{ \begin{pmatrix} w_1 & -\overline{w_2} & ix_1 \\ -w_3 & ix_2 & w_2 \\ ix_3 & \overline{w_3} & -\overline{w_1} \end{pmatrix} : w_1, w_2, w_3 \in \mathbb{C} \text{ and } x_1, x_2, x_3 \in \mathbb{R} \right\}$$

of the indefinite unitary group $U(2, 1)$, where the model of $U(2, 1)$ being used is the subgroup of $GL(3, \mathbb{C})$ that fixes the Hermitian form in 3-variables

$$H(z) = z_1\overline{z_3} + z_3\overline{z_1} + z_2\overline{z_2}.$$

Indeed, writing

$$(3.10) \quad L = -\frac{1}{2}(L_2 - iL_1) \quad \text{and} \quad \varphi = -\frac{1}{2}(3\theta_0^0 + i(\theta_2^1 - \theta_1^2)),$$

we have

$$(3.11) \quad d\zeta = -\gamma \wedge \zeta,$$

where

$$\gamma = \begin{pmatrix} \varphi & -\frac{1}{2}\overline{\zeta_1} & 0 \\ -\frac{1}{2}\zeta_3 & \varphi - \overline{\varphi} & \frac{1}{2}\zeta_1 \\ L\overline{\zeta_1} - \overline{L}\zeta_1 & \frac{1}{2}\overline{\zeta_3} & -\overline{\varphi} \end{pmatrix},$$

as the reader can verify by using the definitions (3.8),(3.10) and the structure equations (3.4). It follows that \mathfrak{J} is integrable. Likewise, the reader may verify that

$$(3.12) \quad d\varphi = \frac{1}{2}\zeta_3 \wedge \overline{\zeta_1} - \frac{1}{4}\zeta_2 \wedge \overline{\zeta_2} - \zeta_1 \wedge \overline{\zeta_3},$$

simply by plugging in the definitions of the involved forms and by using the structure equations (3.4).

Now consider the map

$$\tilde{\psi} : P \rightarrow P \times \mathbb{C}_2 \setminus \{0\}, \quad u \mapsto (u, (0 \ 1))$$

and let $q : P \times \mathbb{C}_2 \setminus \{0\} \rightarrow \mathbb{P}(T_{\mathbb{C}}^*Z^{1,0})$ denote the natural quotient projection induced by (the projectivisation of) ρ . Then $q \circ \tilde{\psi} : P \rightarrow \mathbb{P}(T_{\mathbb{C}}^*Z^{1,0})$ is a submersion onto Y whose fibres are the $\mathrm{CO}(2)$ -orbits. Indeed, let (u, w) be a representative of an element $[v] \in \mathbb{P}(T_{\mathbb{C}}^*Z^{1,0})$ which lies in the complement of $\zeta_1(Z)$. Then using (3.7) it follows that we might transform with the affine part of the right action of $\mathbb{R}_2 \rtimes \mathrm{CO}(2)$ to ensure that w is of the form $(0 \ w_2)$ for some non-zero complex number w_2 . It follows that the element $u \in P$ is mapped onto $[v]$ showing that $q \circ \tilde{\psi}$ is surjective onto Y . Clearly $q \circ \tilde{\psi}$ is smooth and a submersion. Furthermore, suppose the two points $u, u' \in P$ are mapped to the same element of Y . Then, there exists an element $z \rtimes re^{i\phi} \in \mathbb{R}_2 \rtimes \mathrm{CO}(2)$ and a non-zero complex number s so that

$$\rho\left((z \rtimes re^{i\phi})^{-1}\right)(0 \ 1) = (-zr^2e^{-2i\phi} \ e^{-2i\phi}) = (0 \ s)$$

which holds true if and only if $z = 0$. Consequently, there exists a unique diffeomorphism $\psi : P/\mathrm{CO}(2) \rightarrow Y$ making the following diagram commute:

$$\begin{array}{ccc} P & \xrightarrow{\tilde{\psi}} & P \times \mathbb{C}_2 \setminus \{0\} \\ \downarrow \tau & & \downarrow q \\ P/\mathrm{CO}(2) & \xrightarrow{\psi} & Y \end{array}$$

The complex structure on $Y \subset \mathbb{P}(T_{\mathbb{C}}^*Z^{1,0})$ is such that its $(1,0)$ -forms pull back to $P \times \mathbb{C}_2 \setminus \{0\}$ to become linear combinations of the complex-valued 1-forms $\zeta_1, \zeta_2, dw_1, dw_2$, where $w = (w_1 \ w_2) : P \times \mathbb{C}_2 \rightarrow \mathbb{C}_2$ denotes the projection onto the linear factor. Clearly, these forms pull back under $\tilde{\psi}$ to become linear combinations of the forms $\zeta_1, \zeta_2, \zeta_3$, hence ψ is a biholomorphism.

Finally, note that the complex version of the Liouville 1-form on $T_{\mathbb{C}}^*Z^{1,0}$ – whose kernel defines the canonical contact structure on $\mathbb{P}(T_{\mathbb{C}}^*Z^{1,0})$ – pulls back to $P \times \mathbb{C}_2$ to become $w_1\zeta_1 + w_2\zeta_2$. Since

$$\tilde{\psi}^*(w_1\zeta_1 + w_2\zeta_2) = \zeta_2,$$

the claim follows. \square

Remark 3.4. Whereas the definition of the forms ζ_i is a natural consequence of the Lie algebra structure of $\mathrm{CO}(2) \subset \mathbb{R}_2 \rtimes \mathrm{GL}^+(2, \mathbb{R})$, the definition of the form φ in (3.10) is somewhat mysterious at this point. The choice will be clarified during the proof of Theorem 4.9 below.

We will henceforth identify $Y \simeq P/\mathrm{CO}(2)$ and think of τ as the projection map onto Y . Denoting the integrable almost complex structure on Y by J , the first part of the following proposition is therefore clear:

Proposition 3.5. *There exists a unique indefinite Kähler structure on (Y, J) whose Kähler-form Ω_Y satisfies*

$$\tau^* \Omega_Y = -\frac{i}{4} \left(\zeta_1 \wedge \bar{\zeta}_3 + \zeta_3 \wedge \bar{\zeta}_1 + \zeta_2 \wedge \bar{\zeta}_2 \right).$$

Moreover, the indefinite Kähler metric $h_p(\cdot, \cdot) := \Omega_Y(J\cdot, \cdot)$ is Einstein with non-zero scalar curvature.

Proof. The first part of the statement is an immediate consequence of the fact that γ takes values in $\mathfrak{u}(2, 1)$. The skeptical reader might also verify this using the structure equations (3.11). Furthermore, by definition, the associated Kähler metric satisfies

$$\tau^* h = \frac{1}{2} \left(\zeta_1 \circ \bar{\zeta}_3 + \zeta_3 \circ \bar{\zeta}_1 + \zeta_2 \circ \bar{\zeta}_2 \right)$$

and hence the forms $\frac{1}{\sqrt{2}} \zeta_i$ are a unitary coframe for $\tau^* h_p$. In order to verify the Einstein condition it is therefore sufficient that the trace of the curvature form

$$\Gamma = d\gamma + \gamma \wedge \gamma$$

is a non-zero constant (imaginary) multiple of $\tau^* \Omega_Y$. We compute

$$\begin{aligned} 0 &= d^2 \zeta_3 \wedge \zeta_1 \wedge \bar{\zeta}_1 = -d(\gamma_{31} \wedge \zeta_1 + \gamma_{32} \wedge \zeta_2 + \gamma_{33} \wedge \zeta_3) \wedge \zeta_1 \wedge \bar{\zeta}_1 \\ &= \zeta_1 \wedge \bar{\zeta}_1 \wedge \left(dL + \frac{1}{2} \bar{L} \zeta_2 - L\varphi - 2L\bar{\varphi} \right), \end{aligned}$$

where we have used (3.11) and (3.12). It follows that there exist unique complex-valued functions L' and L'' on P such that

$$(3.13) \quad dL = L' \zeta_1 + L'' \bar{\zeta}_1 - \frac{1}{2} \bar{L} \zeta_2 + L\varphi + 2L\bar{\varphi}.$$

Using the structure equations (3.11), (3.12) and (3.13) we compute

$$\Gamma = \frac{1}{4} \begin{pmatrix} \Gamma_{11} & -\zeta_1 \wedge \bar{\zeta}_2 & \zeta_1 \wedge \bar{\zeta}_1 \\ -\zeta_2 \wedge \bar{\zeta}_3 & \Gamma_{22} & \bar{\zeta}_1 \wedge \zeta_2 \\ \zeta_3 \wedge \bar{\zeta}_3 + * & \bar{\zeta}_2 \wedge \zeta_3 & \Gamma_{33} \end{pmatrix},$$

with

$$\begin{aligned} \Gamma_{11} &= \frac{1}{4} \left(\zeta_3 \wedge \bar{\zeta}_1 - \zeta_2 \wedge \bar{\zeta}_2 - 4\zeta_1 \wedge \bar{\zeta}_3 \right), \\ \Gamma_{22} &= \frac{1}{4} \left(-\zeta_1 \wedge \bar{\zeta}_3 - 2\zeta_2 \wedge \bar{\zeta}_2 - \zeta_3 \wedge \bar{\zeta}_1 \right), \\ \Gamma_{33} &= \frac{1}{4} \left(\zeta_1 \wedge \bar{\zeta}_3 - \zeta_2 \wedge \bar{\zeta}_2 - 4\zeta_3 \wedge \bar{\zeta}_1 \right). \end{aligned}$$

and where $*$ = $4(L' + \bar{L}') \zeta_1 \wedge \bar{\zeta}_1$. In particular, we obtain

$$\Gamma_{11} + \Gamma_{22} + \Gamma_{33} = 4i\tau^* \Omega_Y,$$

thus verifying the Einstein property. \square

Remark 3.6. In [15], it is shown how to canonically associate a split-signature anti-self-dual Einstein metric on the total space of a certain rank two affine bundle A fibering over a projective surface (Σ, p) . The indefinite Kähler–Einstein manifold (Y, J, Ω_Y) constructed here may be interpreted as the twistor space of this anti-self-dual Einstein metric.

3.4. The canonical flat case

In this subsection we identify the spaces

$$Y = P/\mathrm{CO}(2) \quad \text{and} \quad Z = P/(\mathbb{R}_2 \rtimes \mathrm{CO}(2))$$

in the case where (Σ, p) is the canonical flat projective structure on the projective 2-sphere. Recall that in this case $P = \mathrm{SL}(3, \mathbb{R})$. The group $\mathrm{SL}(3, \mathbb{R})$ also acts naturally on \mathbb{C}_3 by complexification, that is, by the rule

$$g \cdot (\xi + i\chi) = \xi g^{-1} + i\chi g^{-1}$$

for all $g \in \mathrm{SL}(3, \mathbb{R})$. Clearly, this action descends to define a left action on \mathbb{CP}_2 . However, this action is not transitive, but has two orbits. The first orbit is $\mathbb{RP}_2 \subset \mathbb{CP}_2$, where we think of \mathbb{RP}_2 as those points $[\xi + i\chi] \in \mathbb{CP}_2$ which satisfy $\xi \wedge \chi = 0$, that is, ξ and χ are linearly dependent over \mathbb{R} . Assume therefore $[\varepsilon]$ is an element in the complement $\mathbb{CP}_2 \setminus \mathbb{RP}_2$ of \mathbb{RP}_2 in \mathbb{CP}_2 . Since $\mathrm{SL}(3, \mathbb{R})$ acts transitively on unimodular triples of vectors in \mathbb{R}_3 , we can assume without losing generality that $\varepsilon = (0 \ -i \ 1)$. For $g \in \mathrm{SL}(3, \mathbb{R})$ we write $g = (g_0 \ g_1 \ g_2)$ with $g_i \in \mathbb{R}^3$. We will next determine the stabiliser subgroup of $[\varepsilon]$. A simple computation gives

$$g \cdot \varepsilon = g_0 \wedge (g_1 + ig_2).$$

An elementary calculation shows that $[g \cdot \varepsilon] = [\varepsilon]$ implies that we must have

$$\begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \begin{pmatrix} g_1^2 & -g_1^1 \\ g_2^2 & -g_2^1 \end{pmatrix} \begin{pmatrix} g_0^1 \\ g_0^2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

Since

$$\det g = g_2^0 c_1 - g_1^0 c_2 + g_0^0 \det \begin{pmatrix} g_1^2 & -g_1^1 \\ g_2^2 & -g_2^1 \end{pmatrix} = 1,$$

it follows that $g_0^1 = g_0^2 = 0$. Therefore, the stabiliser subgroup of $[\varepsilon]$ is a subgroup of $\mathbb{R}_2 \rtimes \mathrm{GL}(2, \mathbb{R})$. Writing $a = (a_i^j)$, we obtain

$$(b \rtimes a) \cdot \varepsilon = \det a^{-1} \begin{pmatrix} 0 & -a_1^2 - ia_2^2 & a_1^1 + ia_2^1 \end{pmatrix},$$

from which it follows that $[(b \rtimes a) \cdot \varepsilon] = [\varepsilon]$ if and only if $a_1^1 = a_2^2$ and $a_2^1 + a_1^2 = 0$, that is, $a \in \mathrm{CO}(2)$. Concluding, we have shown

$$\mathrm{SL}(3, \mathbb{R})/(\mathbb{R}_2 \rtimes \mathrm{CO}(2)) \simeq \mathbb{CP}_2 \setminus \mathbb{RP}_2$$

and the projection map is

$$\mu : \mathrm{SL}(3, \mathbb{R}) \rightarrow \mathbb{CP}_2 \setminus \mathbb{RP}_2, \quad (g_0 \ g_1 \ g_2) \mapsto [g_0 \wedge (g_1 + ig_2)],$$

where we use $\mathbb{R}_3 \simeq \Lambda^2(\mathbb{R}^3)$.

Remark 3.7. We have only shown that $Z = \mathrm{SL}(3, \mathbb{R})/(\mathbb{R}_2 \times \mathrm{CO}(2))$ is diffeomorphic to $\mathbb{CP}_2 \setminus \mathbb{RP}_2$. Since Z carries an integrable almost complex structure J , we may ask if (Z, J) is biholomorphic to $\mathbb{CP}_2 \setminus \mathbb{RP}_2$ equipped with the standard complex structure. This is indeed the case, see [37, Prop. 3]. As a consequence of this result one can prove that the conformal connections on the 2-sphere whose (unparametrised) geodesics are the great circles are in one-to-one correspondence with the smooth quadrics in $\mathbb{CP}_2 \setminus \mathbb{RP}_2$, see [37, Cor. 2].

Remark 3.8. In fact [31], if \mathfrak{p} is a projective structure on the 2-sphere, all of whose geodesics are simple closed curves, then Z can be compactified and the compactification is biholomorphic to \mathbb{CP}_2 . This allowed Lebrun and Mason to prove that there is a nontrivial moduli space of such projective structures on the 2-sphere.

We will show next that Y is a submanifold of $F(\mathbb{C}_3)$. Clearly, the action of $\mathrm{SL}(3, \mathbb{R})$ on the space $F(\mathbb{C}_3)$ of complete complex flags is not transitive, there is however an open orbit. Let $F(\mathbb{C}_3)^*$ denote the $\mathrm{SL}(3, \mathbb{R})$ orbit of the flag

$$(\ell, \Pi) = (\mathbb{C}\{\varepsilon_1\}, \mathbb{C}\{\varepsilon_1, \varepsilon_2\}),$$

where

$$\varepsilon_1 = \begin{pmatrix} 0 & -i & 1 \end{pmatrix}, \quad \varepsilon_2 = \begin{pmatrix} 1 & 0 & 0 \end{pmatrix}.$$

We already know that the stabiliser subgroup G_0 of (ℓ, Π) must be a subgroup of $\mathbb{R}_2 \rtimes \mathrm{CO}(2)$. For $b \rtimes a \in \mathbb{R}_2 \rtimes \mathrm{CO}(2)$ we write

$$b \rtimes a = \begin{pmatrix} \frac{1}{x^2+y^2} & b_1 & b_2 \\ 0 & x & y \\ 0 & -y & x \end{pmatrix},$$

with $x^2 + y^2 > 0$. We compute

$$\varepsilon_2 \cdot (b \rtimes a) = \begin{pmatrix} x^2 + y^2 & -xb_1 - yb_2 & -xb_2 + yb_1 \end{pmatrix}$$

which is easily seen to lie in the complex linear span of $\varepsilon_1, \varepsilon_2$ if and only if $b_1 = b_2 = 0$, hence

$$\mathrm{SL}(3, \mathbb{R})/\mathrm{CO}(2) \simeq F(\mathbb{C}_3)^*$$

and the projection map is

$$\tau : \mathrm{SL}(3, \mathbb{R}) \rightarrow F(\mathbb{C}_3) \quad (g_0 \quad g_1 \quad g_2) \mapsto (\mathbb{C}\{\varepsilon_1\}, \mathbb{C}\{\varepsilon_1, \varepsilon_2\}),$$

with

$$\varepsilon_1 = g_0 \wedge (g_1 + ig_2), \quad \varepsilon_2 = g_1 \wedge g_2.$$

Since $F(\mathbb{C}_3)$ is real six-dimensional and since $\dim \mathrm{SL}(3, \mathbb{R}) - \dim \mathrm{CO}(2) = 6$, it follows that $F(\mathbb{C}_3)^*$ is open.

4. The variational equations

By construction, a conformal structure $[g]$ on the oriented projective surface (Σ, \mathfrak{p}) is a section of $Z \rightarrow \Sigma$. Here we will show that every conformal structure $[g]$ admits a natural lift $\widetilde{[g]} : \Sigma \rightarrow Y$. In doing so we recover the functional $\mathcal{E}_{\mathfrak{p}}$ from a different viewpoint, which simplifies the computation of its variational equations. We start with recalling the bundle of complex linear coframes of a Riemann surface.

4.1. The bundle of complex linear coframes

Let Σ be an oriented surface equipped with a conformal structure $[g]$, so that Σ inherits the structure of a Riemann surface whose integrable almost complex structure will be denoted by J . The bundle of complex-linear coframes of $(\Sigma, [g])$ is the $\mathrm{GL}(1, \mathbb{C})$ -subbundle $F_{[g]}^+$ of F^+ consisting of those coframes that are complex-linear with respect to J and the complex structure obtained on \mathbb{R}^2 via the standard identification $\mathbb{R}^2 \simeq \mathbb{C}$. Of course, via the isomorphism $\mathrm{CO}(2) \simeq \mathrm{GL}(1, \mathbb{C})$, we may equivalently think of $F_{[g]}^+$ as consisting of those coframes in F^+ that are angle preserving with respect to $[g]$ and the standard conformal inner product on \mathbb{R}^2 .

Recall that a principal $\mathrm{CO}(2)$ -connection φ on $F_{[g]}^+$ is called *torsion-free* if it satisfies

$$d\omega = -\varphi \wedge \omega,$$

where here we think of the tautological \mathbb{R}^2 -valued 1-form ω on $F_{[g]}^+$ as taking values in \mathbb{C} and the connection taking values in the Lie algebra of $\mathrm{CO}(2) \simeq \mathrm{GL}(1, \mathbb{C})$, that is, \mathbb{C} . The curvature Φ of φ is a $(1,1)$ -form on Σ whose pullback to $F_{[g]}^+$ can be written as

$$d\varphi = R \omega \wedge \bar{\omega}$$

for some unique complex-valued function R on $F_{[g]}^+$. By definition of $F_{[g]}^+$, a complex-valued 1-form on Σ is a $(1,0)$ -form with respect to J if and only if its pullback to $F_{[g]}^+$ is a complex multiple of ω . A consequence of this is the following elementary lemma whose proof we omit:

Lemma 4.1. *A complex-valued function f on $F_{[g]}^+$ represents a section of $K_\Sigma^m \otimes \bar{K}_\Sigma^n$ if and only if there exist complex-valued functions f' and f'' on $F_{[g]}^+$ so that*

$$df = f' \omega + f'' \bar{\omega} + f m \varphi + f n \bar{\varphi}.$$

Remark 4.2. Here $K_\Sigma = T_{\mathbb{C}}^* \Sigma^{1,0}$ denotes the canonical bundle of (Σ, J) , K_Σ^m its m -th tensorial power and \bar{K}_Σ^n the conjugate bundle of the n -th tensorial power of K_Σ . As usual, we let ∇_φ denote the connection induced by φ on $K_\Sigma^m \otimes \bar{K}_\Sigma^n$ and by ∇'_φ its $(1,0)$ -part and by ∇''_φ its $(0,1)$ -part. Of course, if s is the section of $K_\Sigma^m \otimes \bar{K}_\Sigma^n$ represented by f , then $\nabla'_\varphi s$ is represented by f' and $\nabla''_\varphi s$ is represented by f'' .

Theorem 4.1 implies that φ may also be thought of as the connection form of the connection induced by φ on K_Σ^* . Therefore, the first Chern class of K_Σ^* is

$$c_1(K_\Sigma^*) = \left[\frac{i}{2\pi} \Phi \right]$$

and hence if Σ is compact, we obtain

$$(4.1) \quad \int_\Sigma i \Phi = 2\pi \chi(\Sigma),$$

where $\chi(\Sigma)$ denotes the Euler-characteristic of Σ .

4.2. Submanifold theory in the twistor space

We are interested in co-dimension two submanifolds of Z arising as images of sections of $Z \rightarrow \Sigma$. The second order theory of such submanifolds is summarised in the following:

Lemma 4.3. *Let $[g] : \Sigma \rightarrow Z$ be a conformal structure on (Σ, \mathfrak{p}) . Then there exists a lift $\widetilde{[g]} : \Sigma \rightarrow Y$ covering $[g]$ so that the pullback-bundle $p : P'_{[g]} = \widetilde{[g]}^* P \rightarrow \Sigma$ is isomorphic to the $\text{CO}(2)$ -bundle of complex linear coframes $F_{[g]}^+$ of $(\Sigma, [g])$ and so that on $P'_{[g]} \simeq F_{[g]}^+$ we have*

$$\zeta_2 = 2\bar{a}\overline{\zeta_1}, \quad \zeta_3 = k\zeta_1 + 2\bar{q}\overline{\zeta_1},$$

for unique complex-valued functions a, k, q on $P'_{[g]}$.

Proof. First recall that in Theorem 3.3 we have defined

$$\zeta_1 = \theta_0^1 + i\theta_0^2, \quad \zeta_2 = (\theta_1^1 - \theta_2^2) + i(\theta_2^1 + \theta_1^2), \quad \zeta_3 = \theta_1^0 + i\theta_2^0,$$

where $\theta = (\theta_j^i)$ is the Cartan connection of (Σ, \mathfrak{p}) .

Let now $[g] : \Sigma \rightarrow Z$ be a conformal structure on (Σ, \mathfrak{p}) and let $p : P_{[g]} = [g]^* P \rightarrow \Sigma$ denote the pullback of the bundle $\mu : P \rightarrow Z$, that is,

$$P_{[g]} = \{(p, u) \in \Sigma \times P \mid [g](p) = \mu(u)\}.$$

Since $P_{[g]}$ is 6-dimensional, two of the components of θ become linearly dependent when pulled back to $P_{[g]}$. Clearly, these components must be among the 1-forms that are semibasic for μ . Recall that these forms are spanned by ζ_1, ζ_2 and their complex conjugates. However, since $[g]$ is a section of $Z \rightarrow \Sigma$ and since the 1-forms that are semibasic for the projection $\pi : P \rightarrow \Sigma$ are spanned by $\zeta_1, \overline{\zeta_1}$, it follows that $\zeta_1 \wedge \overline{\zeta_1}$ is non-vanishing on $P_{[g]}$. Therefore, on $P_{[g]}$ we have the relation

$$(4.2) \quad \zeta_2 = 2\bar{a}\overline{\zeta_1} + c\zeta_1$$

for unique complex-valued functions a, c . From the equivariance properties of ζ_1, ζ_2 under the $\mathbb{R}_2 \rtimes \text{CO}(2)$ -right action (3.6), we obtain that for all $u \in P_{[g]}$ and $z \rtimes re^{i\phi} \in \mathbb{R}_2 \rtimes \text{CO}(2)$ we have

$$c(u \cdot z \rtimes re^{i\phi}) = r^3 e^{i\phi} c(u) + r^2 z$$

and

$$(4.3) \quad a(u \cdot z \rtimes re^{i\phi}) = r^3 e^{-3i\phi} a(u).$$

It follows that the equation $c = 0$ defines a locus that corresponds to a section $\widetilde{[g]} : \Sigma \rightarrow Y$ covering $[g]$. On the pullback bundle $P'_{[g]} = \widetilde{[g]}^* P$, where

$$P'_{[g]} = \{(p, u) \in \Sigma \times P \mid \widetilde{[g]}(p) = \tau(u)\},$$

we obtain

$$(4.4) \quad \zeta_2 = 2\bar{a}\overline{\zeta_1}.$$

Since $P'_{[g]}$ is 4-dimensional, two of the remaining components of θ become linearly dependent when pulled back to $P'_{[g]}$. Since the 1-forms that are semibasic for the

projection $\tau : P \rightarrow Y$ are spanned by $\zeta_1, \zeta_2, \zeta_3$ and their complex conjugates, it follows as before that

$$(4.5) \quad \zeta_3 = k\zeta_1 + 2\bar{q}\bar{\zeta}_1$$

for unique complex-valued functions k, q .

Now recall that Cartan's bundle $\pi : P \rightarrow \Sigma$ is isomorphic to $F^+ \times \mathbb{R}_2 \rightarrow \Sigma$ equipped with the G -right action (3.1). Therefore, $P_{[g]} \rightarrow \Sigma$ is isomorphic to $F_{[g]}^+ \times \mathbb{R}_2 \rightarrow \Sigma$ and consequently, the bundle $P'_{[g]} \rightarrow \Sigma$ is isomorphic to $F_{[g]}^+ \rightarrow \Sigma$. \square

We also obtain:

Lemma 4.4. *The functions a, k, q and the 1-form φ satisfy the following structure equations on $P'_{[g]} \simeq F_{[g]}^+$*

$$(4.6) \quad da = a'\zeta_1 - q\bar{\zeta}_1 + 2a\varphi - a\bar{\varphi},$$

and

$$dk = k'\zeta_1 + k''\bar{\zeta}_1 + k\varphi + k\bar{\varphi},$$

$$dq = q'\zeta_1 + \frac{1}{2}(\bar{L} + \bar{k}'' - 2\bar{q}a)\bar{\zeta}_1 + 2q\varphi,$$

and

$$(4.7) \quad d\varphi = \left(|a|^2 + \frac{1}{2}k - \bar{k}\right)\zeta_1 \wedge \bar{\zeta}_1$$

for unique complex-valued functions r', k', k'' and q' on $P'_{[g]}$.

Proof. We will only verify the structure equation for a as the other structure equations are derived in an entirely analogous fashion. The structure equations (3.11) and (4.4) gives

$$\begin{aligned} d\bar{\zeta}_2 &= d(2a\zeta_1) = 2da \wedge \zeta_1 + a d\zeta_1 = -\zeta_1 \wedge da + 2a \left(\zeta_1 \wedge \varphi + \frac{1}{2}\bar{\zeta}_1 \wedge \zeta_2 \right) \\ &= -\bar{\zeta}_1 \wedge \bar{\zeta}_3 + \bar{\zeta}_2 \wedge \bar{\varphi} - \bar{\zeta}_2 \wedge \varphi \\ &= 2q\zeta_1 \wedge \bar{\zeta}_1 + 2a\zeta_1 \wedge \bar{\varphi} - 2a\zeta_1 \wedge \varphi, \end{aligned}$$

where we have used (4.5). Equivalently, we obtain

$$0 = \left(da + q\bar{\zeta}_1 - 2a\varphi + a\bar{\varphi} \right) \wedge \zeta_1,$$

which implies (4.6). Finally, the structure equation (5.2) for φ is an immediate consequence of (3.11), (4.4) and (4.5). \square

As we will see next, the functions a, q, k on $P'_{[g]}$ satisfy certain equivariance properties with respect to the $\text{CO}(2)$ -right action on $P'_{[g]}$ and hence represent sections of complex line bundles associated to $p : P'_{[g]} \rightarrow \Sigma$.

Proposition 4.5. *The choice of a conformal structure $[g]$ on (Σ, \mathfrak{p}) determines the following objects:*

- (i) *A torsion-free connection φ on the bundle of complex-linear coframes of $(\Sigma, [g])$;*
- (ii) *A section α of $K_\Sigma^2 \otimes \bar{K}_\Sigma^*$ that is represented by a .*

- (iii) A quadratic differential Q on Σ that is represented by q ;
- (iv) A $(1,1)$ -form κ on Σ that is represented by k .

Moreover, the quadratic differential Q satisfies

$$(4.8) \quad Q = -\nabla''_{\varphi}\alpha.$$

Proof. By construction of $P'_{[g]} \simeq F_{[g]}^+$, a complex-valued 1-form on Σ is a $(1,0)$ -form for the complex structure J induced by $[g]$ and the orientation if and only if its p -pullback to $P'_{[g]}$ is a complex multiple of ζ_1 . Since

$$(R_{re^{i\phi}})^* \zeta_1 = \frac{1}{r^3} e^{i\phi} \zeta_1$$

it follows that the sections of K_{Σ}^2 are in one-to-one correspondence with the complex-valued functions f on $P'_{[g]}$ satisfying

$$(R_{re^{i\phi}})^* f = r^3 e^{-i\phi} r^3 e^{-i\phi} f = r^6 e^{-2i\phi} f.$$

Likewise, it follows that the sections of $K_{\Sigma}^2 \otimes \overline{K}_{\Sigma}^*$ are in one-to-one correspondence with the complex-valued functions f on $P'_{[g]}$ satisfying

$$(R_{re^{i\phi}})^* f = r^3 e^{-i\phi} r^3 e^{-i\phi} \overline{r^{-3} e^{i\phi}} f = r^3 e^{-3i\phi} f$$

and that the sections of $K_{\Sigma} \otimes \overline{K}_{\Sigma}$ are in one-to-one correspondence with the complex valued functions f on $P'_{[g]}$ satisfying

$$(R_{re^{i\phi}})^* f = r^3 e^{-i\phi} r^3 e^{i\phi} f = r^6 f.$$

From (4.5) and (3.9) we obtain that for all $u \in P'_{[g]}$ and $re^{i\phi} \in \text{CO}(2)$

$$k(u \cdot re^{i\phi}) = r^6 k(u),$$

$$q(u \cdot re^{i\phi}) = r^6 e^{-2i\phi} q(u).$$

These equations imply that there exists a unique quadratic differential Q on Σ that is represented by q and a unique $(1,1)$ -form κ on Σ that is represented by k . Furthermore, (4.3) implies that there exists a unique section α of $K_{\Sigma}^2 \otimes \overline{K}_{\Sigma}^*$ that is represented by a .

It follows from the properties (ii) and (iii) of the Cartan connection that φ is a connection 1-form on the $\text{CO}(2)$ -bundle $P'_{[g]} \rightarrow \Sigma$. Its pushforward under the bundle isomorphism $P'_{[g]} \rightarrow F_{[g]}^+$ is then a $\text{CO}(2)$ -connection on $F_{[g]}^+$ which – by abuse of notation – we denote by φ as well. The structure equation (5.2) implies that φ is torsion-free.

Finally, the identity $Q = -\nabla''_{\varphi}\alpha$ is an immediate consequence of the structure equation (4.6) and Theorem 4.1. \square

We call a map $\psi : (M, g) \rightarrow (N, h)$ between two pseudo-Riemannian manifolds *weakly conformal* if $\psi^* h = fg$ for some smooth function f on M . Note that we do not require f to be positive. Two immediate consequences of Theorem 4.5 are:

Corollary 4.6. *Let $[g]$ be a conformal structure on (Σ, p) . Then the lift $\widetilde{[g]} : (\Sigma, [g]) \rightarrow (Y, h_p)$ is weakly conformal if and only if $Q \equiv 0$. Furthermore, the image of $[g] : \Sigma \rightarrow Z$ is a holomorphic curve if and only if $\alpha \equiv 0$. In particular,*

if $[g](\Sigma) \subset Z$ is a holomorphic curve, then $\widetilde{[g]}(\Sigma) \subset Y$ is a holomorphic contact curve.

Remark 4.7. Here we call a holomorphic curve $\Sigma \subset Y$ a *contact curve* if its tangent bundle is contained in the (holomorphic) contact structure of Y .

Proof of Theorem 4.6. By construction, the metric h_p has the property that its pull-back to P is

$$\tau^* h_p = \frac{1}{2} \left(\zeta_1 \circ \overline{\zeta_3} + \zeta_3 \circ \overline{\zeta_1} + \zeta_2 \circ \overline{\zeta_2} \right).$$

Therefore, from (4.4) and (4.5) it follows that

$$(4.9) \quad p^* \left(\widetilde{[g]}^* h_p \right) = \frac{1}{2} \left(4|a|^2 + (k + \bar{k}) \right) \zeta_1 \circ \overline{\zeta_1} + q \zeta_1 \circ \zeta_1 + \bar{q} \overline{\zeta_1} \circ \overline{\zeta_1}.$$

Since a complex-valued 1-form on Σ is a $(1,0)$ -form for the complex structure defined by $[g]$ and the orientation if and only if its p -pullback to $P'_{[g]}$ is a complex multiple of ζ_1 , equation (4.9) implies that $\widetilde{[g]}^* h_p$ is weakly conformal to $[g]$ if and only if q vanishes identically. The first claim follows.

The second part of the claim is an immediate consequence of (4.4) and the characterisation of the complex structures on Z, Y in terms of $\zeta_1, \zeta_2, \zeta_3$ and the characterisation of the holomorphic contact structure in terms of $\zeta_2 = 0$. \square

Remark 4.8. Recall that if $\psi : (\Sigma, [g]) \rightarrow (N, h)$ is a map from a Riemann surface into a (pseudo-)Riemannian manifold, then the $(2,0)$ -part of the pulled back metric $\psi^* h$ is called the *Hopf differential* of ψ . Therefore (4.9) implies that quadratic differential Q is the Hopf differential of $\widetilde{[g]}$.

Theorem 4.5 shows that for every choice of a conformal structure $[g]$ on Σ we obtain a section α of $K_\Sigma^2 \otimes \overline{K}_\Sigma^*$, as well as a connection φ on the principal $\mathrm{GL}(1, \mathbb{C})$ -bundle of complex-linear coframes of $(\Sigma, [g])$. Since $K_\Sigma^2 \otimes \overline{K}_\Sigma^*$ is a subbundle of $T_\mathbb{C}^* \Sigma^2 \otimes T_\mathbb{C} \Sigma$, we may use the canonical real structure of the latter bundle to take the real part of α . Consequently, the real part of α is a 1-form on Σ with values in $\mathrm{End}(T\Sigma)$. We have already encountered an endomorphism valued 1-form $A_{[g]}$ whose properties we discussed in Theorem 2.4. In Theorem 2.6 we have also seen that the choice of a conformal structure $[g]$ on (Σ, p) determines a unique $[g]$ -conformal connection ${}^{[g]}\nabla$ so that ${}^{[g]}\nabla + A_{[g]}$ defines p . On the other hand, φ also induces a $[g]$ -conformal connection on TM which we denote by ∇_φ .

Proposition 4.9. *We have:*

$$(4.10) \quad \nabla_\varphi = {}^{[g]}\nabla,$$

$$(4.11) \quad 2 \operatorname{Re}(\alpha) = A_{[g]},$$

$$(4.12) \quad p^* (|A_{[g]}|_g^2 d\mu_g) = 2i|a|^2 \zeta_1 \wedge \overline{\zeta_1} = -\frac{i}{2} \zeta_2 \wedge \overline{\zeta_2}.$$

Since a $[g]$ -conformal connection ${}^{[g]}\nabla$ has holonomy in $\mathrm{CO}(2)$, it corresponds to a unique torsion-free principal $\mathrm{CO}(2)$ -connection φ on $F_{[g]}^+$, see for instance [6].

Before proving Theorem 4.9 it is helpful to see explicitly how the principal connection φ is constructed from ${}^{[g]}\nabla$. The $[g]$ -conformal connection ${}^{[g]}\nabla$ can be written as

$$(4.13) \quad (g, \beta)\nabla = {}^g\nabla + g \otimes \beta^\sharp - \beta \otimes \text{Id} - \text{Id} \otimes \beta,$$

where $g \in [g]$ and β is a 1-form on M with g -dual vector field β^\sharp . Let $g_{ij} = g_{ji}$ be the unique real-valued functions on F^+ so that $v^*g = g_{ij}\omega^i \otimes \omega^j$. Let $\psi = (\psi_j^i)$ denote the Levi-Civita connection form of g , so that we have the structure equations.

$$\begin{aligned} d\omega^i &= -\psi_j^i \wedge \omega^j, \\ dg_{ij} &= g_{ik}\psi_j^k + g_{kj}\psi_i^k \end{aligned}$$

as well as

$$(4.14) \quad d\psi_j^i + \psi_k^i \wedge \psi_j^k = g_{jk}K_g\omega^i \wedge \omega^k,$$

where the real-valued function K_g on F^+ is (the pullback of) the Gauss curvature of g . Therefore, writing $v^*\beta = b_i\omega^i$ for real-valued functions b_i on F^+ , the connection 1-form of (4.13) is

$$\eta_j^i = \psi_j^i + \left(b_k g^{ki} g_{jl} - \delta_j^i b_l - \delta_l^i b_j\right)\omega^l,$$

where the real-valued functions $g^{ij} = g^{ji}$ on F^+ satisfy $g^{ik}g_{kj} = \delta_j^i$. The equivariance properties of the functions b_i imply that there exist unique real-valued functions b_{ij} on F so that

$$(4.15) \quad db_i = b_j\psi_i^j + b_{ij}\omega^j.$$

From the equivariance properties of the functions g_{ij} it follows that the conditions $g_{11} = g_{22}$ and $g_{12} = 0$ define a reduction of $v : F^+ \rightarrow \Sigma$ to the $\text{CO}(2)$ -subbundle of complex linear coframes of $F_{[g]}^+ \rightarrow \Sigma$ of $(\Sigma, [g])$. On $F_{[g]}^+$ we obtain

$$0 = dg_{12} = g_{11}\psi_2^1 + g_{12}\psi_2^2 + g_{12}\psi_1^1 + g_{22}\psi_1^2 = g_{11}(\psi_2^1 + \psi_1^2)$$

and hence $\psi_1^2 = -\psi_2^1$. Likewise, we have

$$\begin{aligned} 0 &= dg_{11} - dg_{22} = 2(g_{11}\psi_1^1 + g_{12}\psi_1^2) - 2(g_{12}\psi_2^1 + g_{22}\psi_2^2) \\ &= 2g_{11}(\psi_1^1 - \psi_2^2) \end{aligned}$$

so that $\psi_1^1 = \psi_2^2$. Identifying $\mathbb{R}^2 \simeq \mathbb{C}$, we may think of $\omega = (\omega^i)$ as taking values in \mathbb{C} . If we define $\varphi := \frac{1}{2}(\eta_1^1 + \eta_2^2) + \frac{i}{2}(\eta_1^2 - \eta_2^1)$, we obtain

$$(4.16) \quad \varphi = (\psi_1^1 - b_1\omega^1 - b_2\omega^2) + i(\psi_1^2 + b_2\omega^1 - b_1\omega^2)$$

Using this notation the first structure equation can be written in complex form

$$d\omega = -\varphi \wedge \omega,$$

hence φ defines a torsion-free principal $\text{CO}(2)$ -connection on $F_{[g]}^+$.

Proof of Theorem 4.9. Without losing generality, we can assume that the projective structure \mathfrak{p} is defined by ${}^{[g]}\nabla + A_{[g]}$ for some $[g]$ -conformal connection ${}^{[g]}\nabla$ and some 1-form $A_{[g]}$ having all the properties of Theorem 2.4. Recall (3.3) that the

choice of a representative connection $\nabla \in \mathfrak{p}$ gives an identification $P \simeq F^+ \times \mathbb{R}_2$ of Cartan's bundle so that the Cartan connection form becomes

$$(4.17) \quad \theta = \begin{pmatrix} -\frac{1}{3} \operatorname{tr} \eta - \xi \omega & d\xi - \xi \eta - (S\omega)^t - \xi \omega \xi \\ \omega & \eta - \frac{1}{3} \operatorname{tr} \eta + \omega \xi \end{pmatrix}.$$

We will construct Cartan's connection for the representative connection

$$(4.18) \quad (g, \beta) \nabla + A_{[g]} = g \nabla + g \otimes \beta^\# - \beta \otimes \operatorname{Id} - \operatorname{Id} \otimes \beta + A_{[g]}.$$

Let A_{jk}^i denote the real-valued functions on F^+ representing $A_{[g]}$. In particular, we have

$$(4.19) \quad A_{jk}^i = A_{kj}^i \quad \text{and} \quad A_{il}^l = 0.$$

On F^+ the connection form of (4.18) is given by

$$(4.20) \quad \eta_j^i = \psi_j^i + \left(b_k g^{ki} g_{jl} - \delta_j^i b_l - \delta_l^i b_j + A_{jl}^i \right) \omega^l,$$

By definition, the pullback bundle $P_{[g]}$ is the subbundle of $F^+ \times \mathbb{R}_2$ defined by the equations $g_{11} = g_{22}$ and $g_{12} = 0$. Now on $P_{[g]} \simeq F_{[g]}^+ \times \mathbb{R}_2$ we have $\psi_1^2 = -\psi_2^1$ and $\psi_1^1 = \psi_2^2$. Using (4.17), (4.19) and (4.20) we compute

$$\begin{aligned} \zeta_2 &= (\theta_1^1 - \theta_2^2) + i(\theta_2^1 + \theta_1^2) \\ &= \psi_1^1 - \psi_2^2 + (\xi_1 + 2A_{11}^1) \omega^1 + (-\xi_2 - 2A_{22}^2) \omega^2 \\ &\quad + i(\psi_2^1 + \psi_1^2 + (\xi_2 - 2A_{22}^2) \omega^1 + (\xi_1 - 2A_{11}^1) \omega^2) \\ &= 2\bar{a}\bar{\xi}_1 + c\xi_1, \end{aligned}$$

where

$$(4.21) \quad \begin{aligned} a &= A_{11}^1 + iA_{22}^2, \\ c &= \xi_1 + i\xi_2 \end{aligned}$$

and we have used that on $F_{[g]}^+$

$$\delta_{il} A_{jk}^l = \delta_{jl} A_{ik}^l,$$

which follows from Theorem 2.4 (vi). Recall that $P'_{[g]}$ was defined by the equation $c = 0$. Hence on $P'_{[g]} \simeq F_{[g]}^+$ the function ξ vanishes identically. Using this we compute

$$\varphi = -\frac{1}{2} (3\theta_0^0 + i(\theta_2^1 - \theta_1^2)) = \psi_1^1 - b_1 \omega^1 - b_2 \omega^2 + i(\psi_1^2 + b_2 \omega^1 - b_1 \omega^2).$$

This is precisely (4.16). It follows that the connection defined by φ is the same as the induced torsion-free connection on $F_{[g]}^+$ by $^{[g]}\nabla$. This proves (4.10).

Suppose $x = (x^i) : U \rightarrow \mathbb{R}^2$ are local orientation preserving $[g]$ -isothermal coordinates on Σ and write $z = (x^1 + ix^2)$. Applying the exterior derivative to z we obtain a local section $\tilde{z} : U \rightarrow F_{[g]}^+$ so that

$$A_{[g]} = \tilde{z}^* A_{jk}^i dx^j \otimes dx^k \otimes \frac{\partial}{\partial x^i}.$$

By definition of α we have

$$\alpha = \tilde{z}^* a dz \otimes dz \otimes \frac{\partial}{\partial \bar{z}},$$

hence (4.11) is an immediate consequence of (4.21).

Finally, in our coordinates we obtain

$$|A_{[g]}|_g^2 d\mu_g = 4|a|^2 dx^1 \wedge dx^2,$$

so that $p^* (|A_{[g]}|_g^2 d\mu_g) = 2i|a|^2 \zeta_1 \wedge \bar{\zeta}_1 = -\frac{i}{2} \zeta_2 \wedge \bar{\zeta}_2$, as claimed. \square

Note that $A_{[g]}$ vanishes identically if and only if α vanishes identically. Therefore, as an immediate consequence of Theorem 4.9, Theorem 2.6 and Theorem 4.6, we obtain an alternative proof of [37, Theorem 3] (see also [36] for a ‘generalisation’ to higher dimensions):

Theorem 4.10. *A conformal structure $[g]$ on (Σ, \mathfrak{p}) is preserved by a conformal connection defining \mathfrak{p} if and only if the image of $[g] : \Sigma \rightarrow Z$ is a holomorphic curve.*

Remark 4.11. Locally the bundle $Z \rightarrow \Sigma$ always admits sections having holomorphic image and therefore every torsion-free connection on $T\Sigma$ is locally projectively equivalent to a conformal connection (see [37] for additional details).

4.3. Derivation of the variational equations

Applying a technique from [5], we compute the variational equations for the functional $\mathcal{E}_{\mathfrak{p}}$. For a compact domain $\Omega \subset \Sigma$ and a section $[g] : \Sigma \rightarrow Z$ we write

$$\mathcal{E}_{\mathfrak{p}, \Omega}([g]) = \int_{\Omega} |A_{[g]}|_g^2 d\mu_g.$$

Definition 4.12. We say $[g]$ is an $\mathcal{E}_{\mathfrak{p}}$ -critical point or that $[g]$ is *extremal for the projective structure \mathfrak{p}* if for every compact $\Omega \subset \Sigma$ and for every smooth variation $[g]_t : \Sigma \rightarrow Z$ with support in Ω , we have

$$\left. \frac{d}{dt} \right|_{t=0} \mathcal{E}_{\mathfrak{p}, \Omega}([g]_t) = 0.$$

Using this definition we obtain:

Theorem A. *Let (Σ, \mathfrak{p}) be an oriented projective surface. A conformal structure $[g]$ on Σ is extremal for \mathfrak{p} if and only if $\widetilde{[g]} : (\Sigma, [g]) \rightarrow (Y, h_{\mathfrak{p}})$ is weakly conformal.*

Proof. Let $[g] : \Sigma \rightarrow Z$ be a conformal structure and $[g]_t : \Sigma \rightarrow Z$ a smooth variation of $[g]$ with support in some compact set $\Omega \subset \Sigma$ and with $|t| < \varepsilon$. We consider the submanifold of $\Sigma \times P \times (-\varepsilon, \varepsilon)$ defined by

$$P'_{[g]_t} = \left\{ (p, u, t_0) \in \Sigma \times P \times (-\varepsilon, \varepsilon) \mid (p, u) \in P'_{[g]_{t_0}} \right\}$$

and denote by $\iota_{[g]_t} : P'_{[g]_t} \rightarrow \Sigma \times P \times (-\varepsilon, \varepsilon)$ the inclusion map. On $\Sigma \times P \times (-\varepsilon, \varepsilon)$ we define the real-valued 2-form

$$A = -\frac{i}{2} \zeta_2 \wedge \bar{\zeta}_2,$$

where, by abuse of notation, we write ζ_2 for the pullback of ζ_2 to $\Sigma \times P \times (-\varepsilon, \varepsilon)$. Using the structure equations (3.11), we compute

$$(4.22) \quad dA = \frac{i}{2} \left(\zeta_1 \wedge \zeta_3 \wedge \bar{\zeta}_2 - \zeta_2 \wedge \bar{\zeta}_1 \wedge \bar{\zeta}_3 \right).$$

Now Theorem 4.9 implies

$$f(t_0) := \mathcal{E}_{p,\Omega}([g]_t)|_{t=t_0} = \int_{\Omega} \left((\iota_{[g]_t})^* A \right) \Big|_{t=t_0}.$$

Therefore

$$f'(0) = \int_{\Omega} (L_{\partial_t} (\iota_{[g]_t})^* A) \Big|_{t=0} = \int_{\Omega} (\partial_t \lrcorner (\iota_{[g]_t})^* dA) \Big|_{t=0},$$

where L_{∂_t} denotes the Lie-derivative with respect to the vector field ∂_t . It follows from the proof of Theorem 4.3 that on $P'_{[g]_t}$ there exist complex-valued functions a, k, q, B, C such that

$$(4.23) \quad \zeta_2 = 2\bar{a}\bar{\zeta}_1 + B dt \quad \text{and} \quad \zeta_3 = k\zeta_1 + 2\bar{q}\bar{\zeta}_1 + C dt$$

where we now write ζ_i instead of $(\iota_{[g]_t})^* \zeta_i$. Combining (4.22) with (4.23) gives

$$(\iota_{[g]_t})^* dA = i(qB + \bar{q}\bar{B}) dt \wedge \zeta_1 \wedge \bar{\zeta}_1$$

so that

$$(4.24) \quad f'(0) = i \int_{\Omega} (qB + \bar{q}\bar{B}) \zeta_1 \wedge \bar{\zeta}_1 \Big|_{t=0}.$$

Recall that $(R_{r^{e^{i\phi}}})^* \zeta_2 = e^{2i\phi} \zeta_2$ and therefore, by definition, the complex-valued function $B|_{t=0}$ satisfies

$$(R_{r^{e^{i\phi}}})^* (B|_{t=0}) = e^{2i\phi} (B|_{t=0}).$$

Since $(R_{r^{e^{i\phi}}})^* \zeta_1 = r^{-3} e^{i\phi} \zeta_1$ it follows that $B|_{t=0}$ represents a section of $\overline{K}_{\Sigma} \otimes K_{\Sigma}^*$ with support in Ω . Here K_{Σ} denotes the canonical bundle of Σ with respect to the complex structure induced by the orientation and $[g] = [g]_t|_{t=0}$.

It remains to show that every such section in (4.23) with support in Ω can be realised via some variation of $[g]$. We fix a representative metric $g \in [g]$. Let $g_{ij} = g_{ji}$ be the real-valued functions on Cartan's bundle P so that $\pi^* g = g_{ij} \theta_0^i \otimes \theta_0^j$. In particular, from the equivariance properties (ii) of the Cartan connection θ it follows that

$$(R_{b \rtimes a})^* \begin{pmatrix} g_{11} & g_{12} \\ g_{21} & g_{22} \end{pmatrix} = (\det a)^2 a^t \begin{pmatrix} g_{11} & g_{12} \\ g_{21} & g_{22} \end{pmatrix} a.$$

Applying property (iii) of the Cartan connection this implies the existence of unique real-valued functions $g_{ijk} = g_{jik}$ so that

$$(4.25) \quad dg_{ij} = -2g_{ij}\theta_0^0 + g_{kj}\theta_i^k + g_{ik}\theta_j^k + g_{ijk}\theta_0^k.$$

Consider the following conformally invariant functions

$$G = \frac{(g_{11} - g_{22}) + 2ig_{12}}{\sqrt{g_{11}g_{22} - (g_{12})^2}}, \quad H = \frac{g_{11} + g_{22}}{\sqrt{g_{11}g_{22} - (g_{12})^2}}.$$

Translating (4.25) into complex form gives the following structure equation

$$(4.26) \quad dG = G'\zeta_1 + G''\bar{\zeta}_1 + H\zeta_2 + \bar{G}(\bar{\varphi} - \varphi),$$

for unique complex-valued functions G', G'' on P . Clearly, the complex-valued functions G' and G'' can be expressed in terms of the functions g_{ijk} , as $\zeta_1 =$

$\theta_0^1 + i\theta_0^2$. In order to verify (4.26) it is thus sufficient to plug in the definitions of the functions G, H , the definitions of the forms ζ_2, φ and to use

$$dg_{ij} = -2g_{ij}\theta_0^0 + g_{kj}\theta_i^k + g_{ik}\theta_j^k \mod \theta_0^1, \theta_0^2.$$

While this is somewhat tedious, it is straightforward, so we omit the computation.

Fix a section of $\overline{K}_\Sigma \otimes K_\Sigma^*$ with respect to $[g]$ having support in Ω . Such sections are well-known to correspond to endomorphisms of $T\Sigma$ that are trace-free and symmetric with respect to $[g]$. In particular, on P there exist real-valued functions (B_j^i) representing the corresponding endomorphism. The functions satisfy

$$B_i^i = 0 \quad \text{and} \quad g_{ij} B_k^j = g_{kj} B_i^j.$$

as well as the equivariance property

$$(R_{b \rtimes a})^* \begin{pmatrix} B_1^1 & B_2^1 \\ B_1^2 & B_2^2 \end{pmatrix} = a^{-1} \begin{pmatrix} B_1^1 & B_2^1 \\ B_1^2 & B_2^2 \end{pmatrix} a.$$

We define $B = \frac{1}{2}(B_1^1 - B_2^2) + \frac{i}{2}(B_2^1 + B_1^2)$, then B satisfies $(R_{z \rtimes r e^{i\phi}})^* B = e^{2i\phi} B$, hence for sufficiently small t we may vary $[g]$ by defining $[g]_t$ via the zero-locus of the function

$$G_t = G - tBH.$$

Consequently, on

$$P_{[g]_t} = \left\{ (p, u, t_0) \in \Sigma \times P \times (-\varepsilon, \varepsilon) \mid (p, u) \in P_{[g]_{t_0}} \right\}$$

we get

$$\begin{aligned} 0 &= dG_t = dG - dtBH - t d(BH) \\ &= G'\zeta_1 + G''\overline{\zeta_1} + H\zeta_2 + \overline{G}(\overline{\varphi} - \varphi) - dtBH - t d(BH) \\ &= G'\zeta_1 + G''\overline{\zeta_1} + H\zeta_2 + t\overline{B}H(\overline{\varphi} - \varphi) - dtBH - t d(BH) \end{aligned}$$

In particular, if we evaluate this last equation on $P_{[g]_t}|_{t=0}$, we obtain

$$0 = G'\zeta_1 + G''\overline{\zeta_1} + H\zeta_2 - dtBH$$

Since H is non-vanishing on $P_{[g]_t}|_{t=0}$ we must have

$$\zeta_2 = -\frac{G'}{H}\zeta_1 - \frac{G''}{H}\overline{\zeta_1} + Bdt.$$

Since $P'_{[g]_t}$ arises by reducing $P_{[g]_t}$, it follows that on $P'_{[g]_t}|_{t=0}$ we obtain

$$\zeta_2 = -\frac{G''}{H}\overline{\zeta_1} + Bdt,$$

as desired. Finally, we now know that (4.24) must vanish where B is any complex-valued function representing an arbitrary section of $\overline{K}_\Sigma \otimes K_\Sigma^*$ with support in Ω . This is only possible if $q|_{t=0}$ vanishes identically. Applying Theorem 4.6 proves the claim. \square

Remark 4.13. Clearly, if $[g](\Sigma) \subset Z$ is a holomorphic curve, then $\widetilde{[g]} : \Sigma \rightarrow Y$ is weakly conformal. Using the structure equations this can be seen as follows. The image $[g](\Sigma) \subset Z$ is a holomorphic curve if and only if α vanishes identically. However, if α vanishes identically, then so does a and hence (4.6) implies that

q vanishes identically as well. Consequently, every projective structure \mathfrak{p} locally admits a conformal structure $[g]$ so that $\widetilde{[g]}$ is weakly conformal.

We conclude this section by showing that in the compact case $\mathcal{E}_{\mathfrak{p}}([g])$ is – up to a topological constant – just the Dirichlet energy of $\widetilde{[g]} : (\Sigma, [g]) \rightarrow (Y, h_{\mathfrak{p}})$.

Lemma 4.14. *Let (Σ, \mathfrak{p}) be a compact oriented projective surface. Then for every conformal structure $[g] : \Sigma \rightarrow Z$ we have*

$$\int_{\Sigma} |A_{[g]}|_g^2 d\mu_g = 2\pi\chi(\Sigma) + \frac{1}{2} \int_{\Sigma} \mathrm{tr}_g \widetilde{[g]}^* h_{\mathfrak{p}} d\mu_g,$$

where $\chi(\Sigma)$ denotes the Euler-characteristic of Σ .

Proof. Recall from (4.9) that

$$p^* \left(\widetilde{[g]}^* h_{\mathfrak{p}} \right) = \frac{1}{2} \left(4|a|^2 + (k + \bar{k}) \right) \zeta_1 \circ \bar{\zeta}_1 + q \zeta_1 \circ \zeta_1 + \bar{q} \bar{\zeta}_1 \circ \bar{\zeta}_1.$$

Hence we obtain

$$\frac{1}{2} \int_{\Sigma} \mathrm{tr}_g \widetilde{[g]}^* h_{\mathfrak{p}} d\mu_g = \frac{1}{2} \int_{\Sigma} \left(4|a|^2 + (k + \bar{k}) \right) \frac{i}{2} \zeta_1 \wedge \bar{\zeta}_1.$$

Since

$$d\varphi = \left(|a|^2 + \frac{1}{2}k - \bar{k} \right) \zeta_1 \wedge \bar{\zeta}_1,$$

we get

$$\frac{i}{2} (d\varphi - d\bar{\varphi}) = \frac{1}{2} \left(4|a|^2 - (k + \bar{k}) \right) \frac{i}{2} \zeta_1 \wedge \bar{\zeta}_1$$

and thus

$$\begin{aligned} \frac{1}{2} \int_{\Sigma} \mathrm{tr}_g \widetilde{[g]}^* h_{\mathfrak{p}} d\mu_g &= \int_{\Sigma} 2i|a|^2 \zeta_1 \wedge \bar{\zeta}_1 - \int_{\Sigma} \frac{i}{2} (d\varphi - d\bar{\varphi}) \\ &= \int_{\Sigma} |A_{[g]}|_g^2 d\mu_g - 2\pi\chi(\Sigma), \end{aligned}$$

where we have used (I.1) and (4.12). \square

As an obvious consequence of Theorem 4.14 and Theorem 2.4 we have the lower bound:

Theorem B. *Let (Σ, \mathfrak{p}) be a compact oriented projective surface. Then for every conformal structure $[g] : \Sigma \rightarrow Z$ we have*

$$\frac{1}{2} \int_{\Sigma} \mathrm{tr}_g \widetilde{[g]}^* h_{\mathfrak{p}} d\mu_g \geq -2\pi\chi(\Sigma),$$

with equality if and only if \mathfrak{p} is defined by a $[g]$ -conformal connection.

5. Existence of critical points

Clearly, if a projective structure \mathfrak{p} is defined by a $[g]$ -conformal connection, then the conformal structure $[g]$ is a critical point for $\mathcal{E}_{\mathfrak{p}}$ and moreover an absolute minimiser. In this final section we study the projective structures for which $\mathcal{E}_{\mathfrak{p}}$ admits a critical point in some more detail. In particular, we will prove that properly convex projective structures admit critical points.

Recall that the choice of a conformal structure $[g]$ on an oriented projective surface (Σ, \mathfrak{p}) determines a torsion-free principal $\mathrm{CO}(2)$ -connection φ on the bundle $F_{[g]}^+$ of complex linear coframes of $(\Sigma, [g])$ and a section α of $K_\Sigma^2 \otimes \overline{K_\Sigma^*}$. Furthermore, the conformal structure $[g]$ is extremal for $\mathfrak{E}_\mathfrak{p}$ if and only if $\nabla_\varphi'' \alpha = 0$. Conversely, let $(\Sigma, [g])$ be a Riemann surface. Let φ be a torsion-free principal $\mathrm{CO}(2)$ -connection on $F_{[g]}^+$ and α a section of $K_\Sigma^2 \otimes \overline{K_\Sigma^*}$. Then Theorem 4.5, Theorem 4.9 and Theorem A show that the conformal structure $[g]$ is extremal for the projective structure defined by $\nabla_\varphi + 2 \operatorname{Re}(\alpha)$ if and only if $\nabla_\varphi'' \alpha \equiv 0$. Since the curvature of the connection induced by φ on the complex line bundle $E = K_\Sigma^2 \otimes \overline{K_\Sigma^*}$ is a $(1,1)$ -form, standard results imply (see for instance [25, Prop. 1.3.7]) that there exists a unique holomorphic line bundle structure $\bar{\partial}_E$ on E , so that

$$\bar{\partial}_E = \nabla_\varphi''.$$

Hence the variational equation $\nabla_\varphi'' \alpha = 0$ just says that α is holomorphic with respect to $\bar{\partial}_E$. Since the line bundle E has degree

$$\deg(E) = \deg(K_\Sigma^2) - \deg(K_\Sigma^*) = -3 \deg(K_\Sigma^*) = -3\chi(\Sigma),$$

we immediately obtain:

Theorem 5.1. *Suppose \mathfrak{p} is a projective structure on the oriented 2-sphere S^2 admitting an extremal conformal structure $[g]$. Then \mathfrak{p} is defined by a $[g]$ -conformal connection.*

Proof. Suppose $[g]$ is an extremal conformal structure of $\mathfrak{E}_\mathfrak{p}$. From Theorem 2.6 we know that \mathfrak{p} is defined by ${}^{[g]}\nabla + A_{[g]}$ for some $[g]$ -conformal connection ${}^{[g]}\nabla$. Since $\chi(S^2) = 2$, we have $\deg(E) = -6$ and hence the only holomorphic section of E is the zero-section. It follows that α vanishes identically and since by Theorem 4.9 we have $A_{[g]} = 2 \operatorname{Re}(\alpha)$, so does $A_{[g]}$. \square

Remark 5.2. Note that the projectively flat conformal connections on S^2 are classified in [37].

From the Riemann–Roch theorem we know that the space $H^0(\Sigma, E)$ of holomorphic sections of E has dimension

$$\dim_{\mathbb{C}} H^0(\Sigma, E) \geq \deg(E) + 1 - g_\Sigma = 5g_\Sigma - 5,$$

where here g_Σ denotes the genus of Σ . In particular, if Σ has negative Euler-characteristic, then $\dim_{\mathbb{C}} H^0(\Sigma, E)$ will have positive dimension.

5.1. Convex projective structures

Recall that a flat projective surface (Σ, \mathfrak{p}) has the property that Σ can be covered with open subsets, each of which is diffeomorphic onto a subset of \mathbb{RP}^2 in such a way that the geodesics of \mathfrak{p} are mapped onto (segments) of projective lines $\mathbb{RP}^1 \subset \mathbb{RP}^2$. This condition turns out to be equivalent to Σ carrying an atlas modelled on \mathbb{RP}^2 , that is, an atlas whose chart transitions are restrictions of fractional linear transformations. On the universal cover $\tilde{\Sigma}$ of the surface the charts can be adjusted to agree on overlaps, thus defining a *developing map* $\operatorname{dev} : \tilde{\Sigma} \rightarrow \mathbb{RP}^2$, unique up to post-composition with an element of $\mathrm{SL}(3, \mathbb{R})$. In addition, one obtains a

monodromy representation $\rho : \pi_1(\Sigma) \rightarrow \mathrm{SL}(3, \mathbb{R})$ of the fundamental group $\pi_1(\Sigma)$ – well defined up to conjugation – making dev into an equivariant map. A flat projective structure is called *properly convex* if dev is a diffeomorphism onto a subset of \mathbb{RP}^2 which is bounded and convex. If Σ is a compact orientable surface with negative Euler characteristic, then (the conjugacy class of) ‘the’ monodromy representation ρ of a properly convex projective structure is an element in the Hitchin component \mathcal{H}_3 of Σ and conversely every element in \mathcal{H}_3 can be obtained in this way [10].

Motivated by the circle of ideas discussed in the introduction, it is shown in [27] and [34] that on a compact oriented surface Σ of negative Euler characteristic, the convex projective structures are parametrised in terms of pairs $([g], C)$, consisting of a conformal structure $[g]$ and a cubic differential C that is holomorphic with respect to the complex structure induced by $[g]$ and the orientation. Indeed, given a holomorphic cubic differential C on such a Σ , there exists a unique Riemannian metric g in the conformal equivalence class $[g]$, so that

$$(5.1) \quad K_g = -1 + 2|C|_g^2,$$

where K_g denotes the Gauss curvature of g and $|C|_g$ the pointwise norm of C with respect to the Hermitian metric induced by g on the third power of the canonical bundle K_Σ of Σ . Now there exists a unique section α of $K_\Sigma^2 \otimes \overline{K}_\Sigma^*$, so that $\alpha \otimes d\mu_g = C$, where here we think of the area form $d\mu_g$ of g as a section of $K_\Sigma \otimes \overline{K}_\Sigma$. Consequently, we obtain a connection $\nabla = {}^g\nabla + 2\mathrm{Re}(\alpha)$ on $T\Sigma$. The projective structure defined by ∇ is properly convex and conversely every properly convex projective structure arises in this way [27, Theorem 4.1.1, Theorem 4.2.1]. The metric g is known as the *affine metric* or *Blaschke metric*, due to the fact that its pullback to the universal cover $\tilde{\Sigma}$ of Σ can be realised via some immersion $\tilde{\Sigma} \rightarrow \mathbb{A}^3$ as a complete hyperbolic affine 2-sphere in the affine 3-space \mathbb{A}^3 . In particular, (5.1) is known as Wang’s equations in the affine sphere literature [43]. We refer the reader to the survey articles [23], [33] as well as [1] for additional details.

Calling a conformal structure $[g]$ on (Σ, \mathfrak{p}) *closed*, if the associated connection φ on $F_{[g]}^+$ induces a flat connection on $\Lambda^2(T^*\Sigma)$, we obtain a novel characterisation of properly convex projective structures among flat projective structures:

Theorem C. *Let (Σ, \mathfrak{p}) be a compact oriented flat projective surface of negative Euler characteristic. Suppose \mathfrak{p} is properly convex, then the conformal equivalence class of the Blaschke metric is closed and extremal for $\mathcal{E}_\mathfrak{p}$. Conversely, if $\mathcal{E}_\mathfrak{p}$ admits a closed extremal conformal structure $[g]$, then \mathfrak{p} is properly convex and $[g]$ is the conformal equivalence class of the Blaschke metric of \mathfrak{p} .*

Remark 5.3. It would be interesting to know if flat projective surfaces (Σ, \mathfrak{p}) exist for which $\mathcal{E}_\mathfrak{p}$ admits an extremal conformal structure that is not closed.

Proof of Theorem C. Assume \mathfrak{p} is properly convex and let $([g], C)$ be the associated pair. Let g the Blaschke metric satisfying (5.1) and φ the connection on $F_{[g]}^+$ induced by the Levi-Civita connection of g . Recall that ∇_φ denotes the connection induced by φ on TM , hence here we have $\nabla_\varphi = {}^g\nabla$. From [27] we know that \mathfrak{p} is defined by a connection of the form

$$\nabla = {}^g\nabla + 2\mathrm{Re}(\alpha),$$

where α satisfies $\alpha \otimes d\mu_g = C$. A simple computation shows that a torsion-free connection φ on $F_{[g]}^+$ induces a flat connection on $\Lambda^2(T^*\Sigma)$ if and only if ∇_φ has symmetric Ricci tensor. Since here $\nabla_\varphi = {}^g\nabla$ is a Levi-Civita connection, it follows that the conformal structure defined by the Blaschke metric is closed. In addition, since C is holomorphic, we have $\nabla_\varphi'' C = 0$ and furthermore, since $d\mu_g$ is parallel with respect to ${}^g\nabla$, it follows that $\nabla_\varphi'' \alpha$ vanishes identically, thus showing that the conformal structure defined by the Blaschke metric is extremal for \mathcal{E}_p .

Conversely, let (Σ, p) be a compact oriented flat projective surface of negative Euler characteristic. Suppose $[g]$ is a closed and extremal conformal structure for p . We let φ denote the induced connection on $F_{[g]}^+$ and α the corresponding section of $K_\Sigma^2 \otimes \overline{K}_\Sigma^*$. Theorem 4.4 implies that on $P'_{[g]} \simeq F_{[g]}^+$ we have the following structure equations, where we write ω instead of ζ_1

$$(5.2) \quad \begin{aligned} da &= a'\omega - q\overline{\omega} + 2a\varphi - a\overline{\varphi}, \\ dk &= k'\omega + k''\overline{\omega} + k\varphi + k\overline{\varphi}, \\ dq &= q'\omega + \frac{1}{2}(\overline{L} + \overline{k}'' - 2\overline{q}a)\overline{\omega} + 2q\varphi, \\ d\varphi &= \left(|a|^2 + \frac{1}{2}k - \overline{k}\right)\omega \wedge \overline{\omega}. \end{aligned}$$

Since $[g]$ is extremal, we know that Q and hence q vanishes identically. Moreover, recall that p is flat if and only if $L \equiv 0$, hence the third structure equation gives

$$0 = dq = q'\omega + \frac{1}{2}\overline{k}''\overline{\omega}$$

showing that the functions q' and k'' vanish identically as well. Theorem 4.1 implies that $-(\varphi + \overline{\varphi})$ is the connection form of the connection induced by φ on $\Lambda^2(T^*\Sigma)$. Since $[g]$ is closed, the induced connection is flat and hence $d(\varphi + \overline{\varphi})$ must vanish identically. Thus we obtain

$$0 = d(\varphi + \overline{\varphi}) = \frac{3}{2}(\overline{k} - k)\omega \wedge \overline{\omega},$$

showing that k must be real-valued. Note that since k is real-valued, we have

$$0 = d(k - \overline{k}) = k'\omega - \overline{k}'\overline{\omega},$$

so that k' vanishes identically. Finally, we have reduced the structure equations to

$$(5.3) \quad \begin{aligned} da &= a'\omega + 2a\varphi - a\overline{\varphi}, \\ dk &= k\varphi + k\overline{\varphi}, \\ d\varphi &= \left(|a|^2 - \frac{1}{2}k\right)\omega \wedge \overline{\omega}. \end{aligned}$$

The equivariance property of the tautological 1-form ω on $F_{[g]}^+$ gives

$$(R_{re^{i\phi}})^*\omega = \frac{1}{r}e^{i\phi}\omega$$

for all $re^{i\phi} \in \text{CO}(2)$. The function k represents a $(1,1)$ -form κ on Σ which satisfies $v^*\kappa = \frac{i}{2}k\omega \wedge \overline{\omega}$. Consequently, k has the equivariance property $(R_{re^{i\phi}})^*k = r^2k$.

Recall that

$$\int_{\Sigma} \text{id}\varphi = \int_{\Sigma} i \left(|a|^2 - \frac{1}{2}k \right) \omega \wedge \bar{\omega} = 2\pi\chi(\Sigma) < 0,$$

hence k must be positive somewhere. Note that (5.3) shows that the $(1,1)$ -form κ represented by k is parallel with respect to φ . Consequently, k cannot vanish. Since Σ is assumed to be connected, the equivariance property of k implies that the equation $k = 1$ defines a reduction $F_g^+ \subset F_{[g]}^+$ to an $\text{SO}(2)$ -subbundle which is the orthonormal coframe bundle of a unique representative metric $g \in [g]$. On F_g^+ we have

$$0 = dk = \varphi + \bar{\varphi},$$

showing that we may write $\varphi = i\phi$ for a unique 1-form ϕ on F_g^+ . Of course, ϕ is the Levi-Civita connection form of g and hence using $\omega = \omega^1 + i\omega^2$, we obtain the familiar structure equation for the Levi-Civita connection of an oriented Riemannian 2-manifold

$$d\phi = -(-1 + 2|a|^2)\omega^1 \wedge \omega^2.$$

We may define a cubic differential C by setting $C = \alpha \otimes d\mu_g$ and since the pullback to F_g^+ of the area form of g is $\omega^1 \wedge \omega^2$, we conclude the the cubic differential C is holomorphic and represented by the function a . Since

$$d\phi = -K_g \omega^1 \wedge \omega^2,$$

where K_g denotes the Gauss curvature of g , we have

$$K_g = -1 + 2|C|_g^2,$$

where we use that $v^*|C|_g^2 = |c|^2$. It follows that g is the Blaschke metric associated to the pair $([g], C)$ and hence \mathfrak{p} is a properly convex projective structure. \square

5.2. Concluding remarks

Remark 5.4. Let G_0 be a real split simple Lie group and $S(G_0)$ the associated symmetric space. For our purposes we may take $G_0 = \text{SL}(3, \mathbb{R})$ so that $S(G_0) = \text{SL}(3, \mathbb{R})/\text{SO}(3)$, but the following results hold in the more general case. Suppose Σ is a compact oriented surface of negative Euler characteristic and $\rho : \pi_1(\Sigma) \rightarrow G_0$ a representation in the Hitchin component for G_0 . By a theorem of Corlette [11], the choice of a conformal structure $[g]$ on Σ determines a map $\psi : \tilde{\Sigma} \rightarrow S(G_0)$ which is equivariant with respect to ρ and harmonic with respect to the Riemannian metric on $S(G_0)$ and the conformal structure on $\tilde{\Sigma}$ obtained by lifting $[g]$. Furthermore, this map is unique up to post-composition with an isometry of $S(G_0)$. The energy density of the map ψ descends to define a 2-form $e_\rho([g]) d\mu_g$ on Σ and hence one may define an energy functional [12], [29]

$$\mathcal{E}_\rho([g]) = \int_{\Sigma} e_\rho([g]) d\mu_g.$$

The energy $\mathcal{E}_\rho([g])$ turns out to only depend on the diffeotopy class of $[g]$ and thus defines an energy functional on Teichmüller space for every representation ρ in the Hitchin component of G_0 . The Hopf differential of the map ψ yields a holomorphic quadratic differential which descends to Σ as well and it is conjectured [17], [29], that for every representation in the Hitchin component there

exists a unique conformal structure on Σ whose associated Hopf differential vanishes identically. For such a conformal structure the mapping ψ is harmonic and conformal, hence minimal. In [30] Labourie proves the existence of a unique ρ -equivariant minimal mapping $\psi : \tilde{\Sigma} \rightarrow S(G_0)$ in the case where G_0 has rank two (the case $G_0 = \mathrm{SL}(3, \mathbb{R})$ was treated previously in [27]). Labourie also shows the existence of such a mapping without any assumption on the rank of G_0 in [29]. Moreover, in [30], the energy bound

$$\mathcal{E}_\rho([g]) \geq -2\pi\chi(\Sigma)$$

is obtained, with equality if and only if ρ is a *Fuchsian representation*.

Given our results it is natural to expect a relation between \mathcal{E}_ρ and our functional \mathcal{E}_p , where ρ is an element in the $\mathrm{SL}(3, \mathbb{R})$ Hitchin component and p denotes its associated properly convex projective structure. However, relating the representation ρ to its associated projective structure p in a way that would allow to establish the expected relation proves to be quite difficult. This may be investigated elsewhere.

Remark 5.5. Although we are currently unable to prove this, the previous remark suggests that in the case of a properly convex compact oriented projective surface (Σ, p) of negative Euler characteristic, the conformal equivalence class of the Blaschke metric is in fact the unique critical point of \mathcal{E}_p . As a partial result towards this claim, it is shown in [41] that if a properly convex compact oriented projective surface (Σ, p) of negative Euler characteristic admits a compatible Weyl connection, then p arises from a hyperbolic metric.

Remark 5.6. In [38], it is shown that for a compact oriented projective surface (Σ, p) of negative Euler characteristic the functional \mathcal{E}_p admits at most one absolute minimiser $[g]$ (i.e. a conformal structure $[g]$ such that $\mathcal{E}_p([g]) = 0$).

Remark 5.7. In [39], the author shows that properly convex projective surfaces arise from torsion-free connections on $T\Sigma$ that admit an interpretation as Lagrangian minimal surfaces. Some of their properties are studied in [40]. It would be interesting to relate these minimal Lagrangian surfaces to the minimal mapping ψ constructed in [27].

Remark 5.8. We have seen that oriented projective structures admitting extremal conformal structures arise from pairs (φ, α) on a Riemann surface $(\Sigma, [g])$, where α satisfies $\nabla_\varphi''\alpha \equiv 0$. The torsion-free connection φ on $F_{[g]}^+$ induces a holomorphic line bundle structure $\bar{\partial}_E$ on $E = K_\Sigma^2 \otimes \overline{K_\Sigma^*}$ and conversely, it is easy to see that for every choice of a holomorphic line bundle structure $\bar{\partial}_E$ on E there exists a unique torsion-free connection φ on $F_{[g]}^+$ inducing $\bar{\partial}_E$. Hence we may equivalently describe these projective structures in terms of a pair $(\bar{\partial}_E, \alpha)$ satisfying $\bar{\partial}_E\alpha \equiv 0$.

Remark 5.9. The so-called *naive Einstein* affine hypersurface structures introduced in [16] also provide examples of projective surfaces admitting an extremal conformal structure.

Appendix I. A Gauss–Bonnet type identity

As a by-product of our considerations, we obtain a Gauss–Bonnet type identity:

Theorem I.1. *Let (Σ, \mathfrak{p}) a compact oriented projective surface. Then for every section $s : \Sigma \rightarrow (Y, \Omega_{\mathfrak{p}})$ we have*

$$(I.1) \quad \int_{\Sigma} s^* \Omega_{\mathfrak{p}} = 2\pi \chi(\Sigma).$$

Proof. Since $\pi : Y \rightarrow \Sigma$ admits smooth global sections, it follows that $\pi^* : H^k(\Sigma) \rightarrow H^k(Y)$ is injective. Note that by construction the fibres of the bundle $\pi : Y \rightarrow \Sigma$ are diffeomorphic to $(\mathbb{R}_2 \rtimes \mathrm{GL}^+(2, \mathbb{R})) / \mathrm{CO}(2)$ and hence diffeomorphic to $\mathbb{R}_2 \times D^2$. In particular, the fibre is contractible, thus we have $H^2(Y) \simeq H^2(\Sigma) \simeq \mathbb{R}$ showing that $\pi^* : H^2(\Sigma) \rightarrow H^2(Y)$ is an isomorphism. It follows that any two sections of $Y \rightarrow \Sigma$ induce the same map on the second de Rham cohomology groups. It is therefore sufficient to construct a section $s : \Sigma \rightarrow Y$ for which (I.1) holds. From the proof of the Theorem 4.3 we know that for every conformal structure $[g] : \Sigma \rightarrow Z$ there exists a lift $\widetilde{[g]} : \Sigma \rightarrow Y$ so that on the pullback bundle $P'_{[g]}$ we have

$$\zeta_2 = 2\bar{a} \bar{\zeta}_1, \quad \zeta_3 = k\zeta_1 + 2\bar{q} \bar{\zeta}_1,$$

Since

$$\tau^* \Omega_{\mathfrak{p}} = -\frac{i}{4} \left(\zeta_1 \wedge \bar{\zeta}_3 + \zeta_3 \wedge \bar{\zeta}_1 + \zeta_2 \wedge \bar{\zeta}_2 \right),$$

computing as in Theorem 4.14 and using the above identities for ζ_2, ζ_3 gives

$$\int_{\Sigma} \widetilde{[g]}^* \Omega_{\mathfrak{p}} = -\frac{i}{4} \int_{\Sigma} \left(k + \bar{k} - 4|a|^2 \right) \zeta_1 \wedge \bar{\zeta}_1 = \frac{i}{2} \int_{\Sigma} d\varphi - d\bar{\varphi} = 2\pi \chi(\Sigma).$$

□

References

- [1] **D. Baraglia**, G_2 geometry and integrable systems, PhD thesis, 2010. arXiv:1002.1767 32
- [2] **Y. Benoist, D. Hulín**, Cubic differentials and finite volume convex projective surfaces, *Geom. Topol.* **17** (2013), 595–620. 4
- [3] **A. L. Besse**, *Einstein manifolds*, *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)* **10**, Springer-Verlag, Berlin, 1987. 7
- [4] **R. Bryant, M. Dunajski, M. Eastwood**, Metrisability of two-dimensional projective structures, *J. Differential Geom.* **83** (2009), 465–499. arXiv:0801.0300 10
- [5] **R. L. Bryant**, A duality theorem for Willmore surfaces, *J. Differential Geom.* **20** (1984), 23–53. 27
- [6] **R. L. Bryant**, Classical, exceptional, and exotic holonomies: a status report, in *Actes de la Table Ronde de Géométrie Différentielle (Luminy, 1992)*, *Sémin. Congr.* **1**, Soc. Math. France, Paris, 1996, pp. 93–165. 24
- [7] **A. Čap, A. R. Gover, H. R. Macbeth**, Einstein metrics in projective geometry, *Geom. Dedicata* **168** (2014), 235–244. arXiv:1207.0128 10
- [8] **A. Čap, J. Slovák**, *Parabolic geometries. I, Mathematical Surveys and Monographs* **154**, American Mathematical Society, Providence, RI, 2009, Background and general theory. 11
- [9] **E. Cartan**, Sur les variétés à connexion projective, *Bull. Soc. Math. France* **52** (1924), 205–241. 11
- [10] **S. Choi, W. M. Goldman**, Convex real projective structures on closed surfaces are closed, *Proc. Amer. Math. Soc.* **118** (1993), 657–661. 3, 32
- [11] **K. Corlette**, Flat G -bundles with canonical metrics, *J. Differential Geom.* **28** (1988), 361–382. 34
- [12] **S. K. Donaldson**, Twisted harmonic maps and the self-duality equations, *Proc. London Math. Soc. (3)* **55** (1987), 127–131. 4, 34

- [13] **M. Dubois-Violette**, Structures complexes au-dessus des variétés, applications, in *Mathematics and physics (Paris, 1979/1982)*, *Progr. Math.* **37**, Birkhäuser Boston, Boston, MA, 1983, pp. 1–42. 2, 13
- [14] **D. Dumas, M. Wolf**, Polynomial cubic differentials and convex polygons in the projective plane, *Geom. Funct. Anal.* **25** (2015), 1734–1798. arXiv:1407.8149 4
- [15] **M. Dunajski, T. Mettler**, Gauge theory on projective surfaces and anti-self-dual Einstein metrics in dimension four, *J. Geom. Anal.* **28** (2018), 2780–2811. arXiv:1509.04276 18
- [16] **D. J. F. Fox**, Einstein-like geometric structures on surfaces, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* (5) **12** (2013), 499–585. arXiv:1011.5723 35
- [17] **W. M. Goldman, R. A. Wentworth**, Energy of twisted harmonic maps of Riemann surfaces, in *In the tradition of Ahlfors-Bers. IV, Contemp. Math.* **432**, Amer. Math. Soc., Providence, RI, 2007, pp. 45–61. arXiv:math/0506212 34
- [18] **O. Guichard, A. Wienhard**, Anosov representations: domains of discontinuity and applications, *Invent. Math.* **190** (2012), 357–438. arXiv:1108.0733 3
- [19] **N. Hitchin**, Complex manifolds and Einstein’s equations, in *Twistor geometry and nonlinear systems, Lecture Notes in Math.* **970**, Springer, Berlin, 1982, pp. 73–99. 2, 13
- [20] **N. Hitchin**, The self-duality equations on a Riemann surface, *Proc. London Math. Soc.* (3) **55** (1987), 59–126. 3
- [21] **N. Hitchin**, Lie groups and Teichmüller space, *Topology* **31** (1992), 449–473. 3
- [22] **N. Hitchin**, Higgs bundles and diffeomorphism groups, in *Surveys in differential geometry 2016. Advances in geometry and mathematical physics, Surv. Differ. Geom.* **21**, Int. Press, Somerville, MA, 2016, pp. 139–163. arXiv:1501.04989 3
- [23] **I. Kim, A. Papadopoulos**, Convex real projective structures and Hilbert metrics, in *Handbook of Hilbert geometry, IRMA Lect. Math. Theor. Phys.* **22**, Eur. Math. Soc., Zürich, 2014, pp. 307–338. arXiv:1406.7079 32
- [24] **S. Kobayashi**, *Transformation groups in differential geometry*, Springer-Verlag, New York, 1972, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 70. 1
- [25] **S. Kobayashi**, *Differential geometry of complex vector bundles*, *Publications of the Mathematical Society of Japan* **15**, Princeton University Press, Princeton, NJ; Princeton University Press, Princeton, NJ, 1987, Kanô Memorial Lectures, 5. 31
- [26] **S. Kobayashi, T. Nagano**, On projective connections, *J. Math. Mech.* **13** (1964), 215–235. 11
- [27] **F. Labourie**, Flat projective structures on surfaces and cubic holomorphic differentials, *Pure Appl. Math. Q.* **3** (2007), 1057–1099. arXiv:math/0611250 4, 32, 35
- [28] **F. Labourie**, Cross ratios, surface groups, $\mathrm{PSL}(n, \mathbf{R})$ and diffeomorphisms of the circle, *Publ. Math. Inst. Hautes Études Sci.* (2007), 139–213. 3
- [29] **F. Labourie**, Cross ratios, Anosov representations and the energy functional on Teichmüller space, *Ann. Sci. Éc. Norm. Supér.* (4) **41** (2008), 437–469. 4, 34, 35
- [30] **F. Labourie**, Cyclic surfaces and Hitchin components in rank 2, *Ann. of Math.* (2) **185** (2017), 1–58. arXiv:1406.4637 4, 35
- [31] **C. Lebrun, L. J. Mason**, Zoll manifolds and complex surfaces, *J. Differential Geom.* **61** (2002), 453–535. arXiv:math/0211021 19
- [32] **R. Liouville**, Sur une classe d’équations différentielles, parmi lesquelles, en particulier, toutes celles des lignes géodésiques se trouvent comprises., *C. R. Acad. Sci., Paris* **105** (1888), 1062–1064. 13
- [33] **J. Loftin**, Survey on affine spheres, in *Handbook of geometric analysis, No. 2, Adv. Lect. Math. (ALM)* **13**, Int. Press, Somerville, MA, 2010, pp. 161–191. arXiv:0809.1186 32
- [34] **J. C. Loftin**, Affine spheres and convex \mathbb{RP}^n -manifolds, *Amer. J. Math.* **123** (2001), 255–274. 4, 32
- [35] **V. S. Matveev, A. Trautman**, A Criterion for Compatibility of Conformal and Projective Structures, *Comm. Math. Phys.* **329** (2014), 821–825. arXiv:1302.2329 2, 8
- [36] **T. Mettler**, Reduction of β -integrable 2-Segre structures, *Comm. Anal. Geom.* **21** (2013), 331–353. arXiv:1110.3279 27
- [37] **T. Mettler**, Weyl metrisability of two-dimensional projective structures, *Math. Proc. Cambridge Philos. Soc.* **156** (2014), 99–113. arXiv:0910.2618 19, 27, 31

- [38] **T. Mettler**, Geodesic rigidity of conformal connections on surfaces, *Math. Z.* **281** (2015), 379–393. arXiv:1410.8501 35
- [39] **T. Mettler**, Minimal Lagrangian connections on compact surfaces, *Adv. Math.* **354** (2019), Article ID 106747, 36 p. arXiv:1609.08033 35
- [40] **T. Mettler, G. Paternain**, Holomorphic differentials, thermostats and Anosov flows, 2019, pp. 553–580. arXiv:1706.03554 35
- [41] **T. Mettler, G. P. Paternain**, Convex projective surfaces with compatible Weyl connection are hyperbolic, 2018, to appear in *Anal. PDE*. arXiv:1804.04616 35
- [42] **N. R. O’Brian, J. H. Rawnsley**, Twistor spaces, *Ann. Global Anal. Geom.* **3** (1985), 29–58. 2, 13
- [43] **C. P. Wang**, Some examples of complete hyperbolic affine 2-spheres in \mathbf{R}^3 , in *Global differential geometry and global analysis (Berlin, 1990)*, *Lecture Notes in Math.* **1481**, Springer, Berlin, 1991, pp. 271–280. arXiv:1504.02551 32
- [44] **H. Weyl**, Zur Infinitesimalgeometrie: Einordnung der projektiven und der konformen Auffassung., *Nachr. Ges. Wiss. Göttingen, Math.-Phys. Kl.* **1921** (1921), 99–112. 2, 6

INSTITUT FÜR MATHEMATIK, GOETHE-UNIVERSITÄT FRANKFURT, FRANKFURT AM MAIN,
GERMANY

Email address: mettler@math.uni-frankfurt.de, mettler@math.ch